

BAX Polyclonal antibody

Catalog Number: 50599-2-Ig

Featured Product

2885 Publications

Basic Information

Catalog Number:

50599-2-Ig

Size:

800 µg/ml

Source:

Rabbit

Isotype:

IgG

Immunogen Catalog Number:

AG0576

GenBank Accession Number:

BC014175

GeneID (NCBI):

581

UNIPROT ID:

Q07812

Full Name:

BCL2-associated X protein

Calculated MW:

21 kDa

Observed MW:

21 kDa

Purification Method:

Protein A purification

Recommended Dilutions:

WB 1:2000-1:16000

IP 0.5-4.0 µg for 1.0-3.0 mg of total protein lysate

IHC 1:1000-1:4000

Applications

Tested Applications:

FC, IHC, IP, WB, ELISA

Cited Applications:

ChIP, CoIP, ELISA, FC, IHC, IP, WB

Species Specificity:

human, mouse, rat

Cited Species:

human, goat, chicken, rat, sheep, mouse, rabbit, zebrafish, hamster, pig

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB: HEK-293 cells, PC-12 cells, HT-1080 cells, HeLa cells, COLO 320 cells, RAW 264.7 cells, NIH/3T3 cells, Raji cells, mouse kidney tissue, rat kidney tissue, mouse heart tissue, rat colon tissue, rat lung tissue, rat brain tissue

IP: Raji cells, mouse heart tissue

IHC: mouse lung tissue, mouse kidney tissue

Background Information

BAX, also named as BCL2L4, is a pro-apoptotic member of the Bcl-2 protein family, which plays a pivotal role in controlling cell life and death. Bax largely localizes to the cytoplasm of healthy cells, but accumulates on the outer mitochondrial membrane upon apoptosis induction (PMID: 9108035). BAX can commit a cell to apoptosis by translocation from the cytosol to the mitochondria and permeabilization of the outer mitochondrial membrane, which leads to the release of cytochrome c from mitochondria (PMID: 21763611). The expression of BAX is upregulated by the tumor suppressor protein p53, and BAX has been shown to be involved in p53-mediated apoptosis (PMID: 8183579).

Notable Publications

Author	Pubmed ID	Journal	Application
Xiaoxuan Zhai	36184541	Int Heart J	WB
Xin Shen	36184549	Int Heart J	WB
Ling Guo	34867437	Front Physiol	WB

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


For technical support and original validation data for this product please contact:

T: 4006900926


E: Proteintech-CN@ptglab.comW: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 50599-2-Ig (BAX antibody) at dilution of 1:8000 incubated at room temperature for 1.5 hours.




WB result of BAX antibody (50599-2-Ig, 1:5000) with si-Control and si-BAX transfected HeLa cells.




Immunohistochemical analysis of paraffin-embedded mouse lung tissue slide using 50599-2-Ig (BAX antibody) at dilution of 1:2000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffin-embedded mouse lung tissue slide using 50599-2-Ig (BAX antibody) at dilution of 1:2000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



IP result of anti-BAX (IP:50599-2-Ig, 4ug; Detection:50599-2-Ig 1:1000) with Raji cells lysate 3160 ug.



1X10⁶ Ramos cells were intracellularly stained with 0.4 ug Anti-Human BAX (50599-2-Ig) and CoraLite®488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L) at dilution 1:1000 (red), or 0.4 ug Control Antibody. Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).