

STAT4 Polyclonal antibody

Catalog Number: 51070-2-AP

2 Publications

Basic Information

Catalog Number:

51070-2-AP

Size:

500 µg/ml

Source:

Rabbit

Isotype:

IgG

GenBank Accession Number:

BC031212

GeneID (NCBI):

6775

UNIPROT ID:

Q14765

Full Name:

signal transducer and activator of transcription 4

Calculated MW:

748 aa, 86 kDa

Observed MW:

86 kDa

Purification Method:

Antigen affinity purification

Recommended Dilutions:

WB 1:2000-1:16000

Applications

Tested Applications:

WB, ELISA

Cited Applications:

WB

Species Specificity:

human, mouse, rat

Cited Species:

human, mouse

Positive Controls:

WB: A549 cells, HeLa cells, mouse testis tissue, rat testis tissue

Background Information

The JAK/STAT pathway is an extensive signaling pathway downstream of cytokine receptors. STATs are cytosolic proteins with a common structure consisting of an N-terminal oligomerization domain, which favors formation of STAT dimers, followed by a DNA-binding domain and a C-terminal SRC homology-2 (SH2) domain, which is involved in association between STATs and receptors [PMID:22383755]. Signal Transducer and Activator of Transcription 4 (STAT4) is a transcription factor that is activated by IL-12 signaling and promotes Th1-cell differentiation and IFN- γ production [PMID:21998209].

Notable Publications

Author	Pubmed ID	Journal	Application
James P Hewitson	32321759	J Immunol	WB
Ke Xiang	38064840	Food Chem	WB

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


For technical support and original validation data for this product please contact:

T: 4006900926

E: Proteintech-CN@ptglab.comW: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 51070-2-AP (STAT4 antibody) at dilution of 1:8000 incubated at room temperature for 1.5 hours.