For Research Use Only

Adenosine A1 Receptor Polyclonal antibody



Catalog Number:55026-1-AP


8 Publications


Basic Information	Catalog Number: 55026-1-AP	GenBank Accession Number: NM_000674		Purification Method: Antigen affinity purification					
	Size:	GeneID (NCBI):		Recommended Dilutions: WB 1:500-1:1000					
	1000 µg/ml								
	Source:UNIPROT ID:RabbitP30542		IHC 1:50-1:500						
	Isotype: Full Name: IgG adenosine A1 receptor								
	Calculated MW: 37 kDa Observed MW: 37 kDa								
Applications	Tested Applications:		Positive Controls:						
			WB:SH-SY	Y5Y cells,					
	Cited Applications: IHC : mouse cerebellum tissue, mouse br WB, IF, IHC : mouse totications:								
	May in find mouse testis tissue Species Specificity: mouse testis tissue human, mouse Cited Species: human, rat, mouse Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0								
					Background Information		portant mediator of e	thanol intoxication and	d by G proteins which inhibit adenylyl d exerts some of its effects via ADORA1
					Notable Publications	Author	Pubmed ID	Journal	Application
Tina M Burke	26378246	Sci Transl Med	WB						
Troy A Hackett	30315630	Anat Rec (Hoboke	n) IHC,IF						
Sen Chen	35468537	Bioorg Med Chem	WB						
	Storage:								

For technical support and original validation data for this product please contact:T: 4006900926E: Proteintech-CN@ptglab.comW: ptgcn.com


This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data





Immunohistochemical analysis of paraffinembedded mouse cerebellum tissue slide using 55026-1-AP (Adenosine A1 Receptor antibody at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded mouse cerebellum tissue slide using 55026-1-AP (Adenosine A1 Receptor antibody at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).

SH-SY5Y cells were subjected to SDS PAGE followed by western blot with 55026-1-AP (Adenosine A1 Receptor antibody) at dilution of 1:500 incubated at room temperature for 1.5 hours.