For Research Use Only

POU4F2 Polyclonal antibody

Catalog Number:55042-1-AP 3 Publications


Antibodies | ELISA kits | Proteins www.ptglab.com

| Basic Information | Catalog Number: GenBank Accession Numb 55042-1-AP NM_004575 | | on Number: | r: Purification Method: Antigen affinity purification | |
|------------------------|--|--------------------------|--------------------|--|--|
| | Size: 280 µg/ml | GenelD (NCBI): 5458 | | Recommended Dilutions: WB 1:500-1:1000 | |
| | Source: UNIPROT ID: Rabbit Q12837 | | | | |
| | Isotype:Full Name:IgGPOU class 4 homeobox 2 | | eobox 2 | | |
| | | Calculated MW: 43 kDa | | | |
| | Observed MW: 40-44 kDa | | | | |
| Applications | Tested Applications: Positive Controls: WB, ELISA | | | | |
| | WB: LO2 cells, mouse eye tissue, mouse brain tissue Cited Applications: IF, WB | | | | |
| | Species Specificity: human, mouse | | | | |
| | Cited Species: human, mouse | | | | |
| Background Information | POU4F2,also named as BRN3.2, BRN3B and Brn-3b, belongs to the POU transcription factor family and Class-4 subfamily. It is a transcription factor. POU4F2 may play a role in determining or maintaining the identities of a small subset of visual system neurons. The antibody is specific to POU4F2. | | | | |
| Notable Publications | Author | Pubmed ID J | ournal | Application | |
| | Zuberwasim Sayyad | 34023378 N | leurochem Int | WB, IF | |
| | Deliang Zhu | 32131893 S | Stem Cell Res Ther | | |
| | Zuberwasim Sayyad | 29203899 S | Sci Rep | WB,IF | |
| Storage | Storage: Store at -20°C. Stable for one year after shipment. Storage Buffer: PBS with 0.02% sodium azide and 50% glycerol pH 7.3. Aliquoting is unnecessary for -20°C storage | | | | |

For technical support and original validation data for this product please contact:T: 4006900926E: Proteintech-CN@ptglab.comW: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



LO2 cells were subjected to SDS PAGE followed by western blot with 55042-1-AP (POU4F2 antibody) at dilution of 1:500 incubated at room temperature for 1.5 hours.