For Research Use Only

Prohibitin Monoclonal antibody

Catalog Number:60092-1-lg 2 Publications



Basic Information

Catalog Number: 60092-1-lg

Size: 900 μg/ml Source: Mouse Isotype: lgG1

Immunogen Catalog Number:

AG1240

Calculated MW: 30 kDa

Observed MW: 30 kDa

BC013401

GeneID (NCBI):

UNIPROT ID:

Full Name:

prohibitin

P35232

GenBank Accession Number:

Purification Method:

Protein A purification

CloneNo.: 4D3G5

Recommended Dilutions:

WB 1:500-1:2000 IHC 1:50-1:500

Applications

Tested Applications: IHC, WB, ELISA Cited Applications:

IF. WB

Species Specificity: human, pig, rat **Cited Species:** human, mouse

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB: human heart tissue, HeLa cells, human placenta tissue, pig heart tissue, rat heart tissue

IHC: human liver tissue, human kidney tissue, human heart tissue

Background Information

Prohibitin, also known as PHB, a member of the Band-7 family of proteins, is widely distributed in bacteria, plants, yeast, protozoa and mammals and is important in cell proliferation, differentiation and apoptosis (PMID: 20840588). This protein localizes to the inner membrane of mitochondria, where it acts as a chaperone protein, and is also found in the nucleus, where it negatively regulates transcription (PMID: 18558096). Recent study confirmed that the expression and distribution of PHB, which is a nuclear matrix protein, affect the apoptosis of HaCaT cells and its colocalization with specific gene products connected with cell apoptosis (PMID: 24402549).

Notable Publications

| Author | Pubmed ID | Journal | Application |
|-------------|-----------|---------------------------------|-------------|
| Yating Peng | 35378837 | Int J Chron Obstruct Pulmon Dis | WB |
| Li-Li Ma | 29100316 | Oncotarget | IF |

Storage


Storage:

Store at -20°C. Stable for one year after shipment.


PBS with 0.1% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 60092-1-1g (Prohibitin antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded human liver tissue slide using 60092-1-Ig (Prohibitin antibody) at dilution of 1:200 (under 10x lens).



Immunohistochemical analysis of paraffinembedded human liver tissue slide using 60092-1-Ig (Prohibitin antibody) at dilution of 1:200 (under 40x lens).