For Research Use Only

## WTAP Monoclonal antibody

Catalog Number:60188-1-lg Featured Product

68 Publications




Basic Information	Catalog Number: 60188-1-lg	GenBank Accession Number: BC000383		Purification Method: Protein A purification
	Size:	GenelD (NCBI):		CloneNo.:
	1000 µ g/ml 9589		4A10G9	
	Source: UNIPROT ID: Mouse Q15007		Recommended Dilutions: WB 1:5000-1:50000	
	lsotype: IgG1	Full Name: Wilms tum	or 1 associated protein	IP 0.5-4.0 ug for 1.0-3.0 mg of total protein lysate
	Immunogen Catalog Number: Calculated MW:   AG0259 45 kDa   Observed MW: 50 kDa		MW:	IHC 1:500-1:2000 IF 1:50-1:500
			1W:	
Applications	Tested Applications:		Positive Controls:	
		IF/ICC, IHC, IP, WB, ELISA WB : MCF-7		ells, HeLa cells, HEK-293 cells, HepG2
	Cited Applications: WB, IP, IF, FC, RIP, IHC	cells, LNCaP cells, 4T1 ce		cells, Jurkat cells, HSC-T6 cells, NIH/3 lls
	Species Specificity: human, mouse, rat	IP : HeLa cel		s,
	Cited Species:	IHC : human colon tissue,		colon tissue,
	human, rat, mouse, drosophila	IF : HeLa cells,		S,
	Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0			
Background Information	WTAP, also named as KIAA0105 and hFL(2)D, is previously identified as a protein associated with Wilms' tumor-1 (WT-1) protein that is essential for the development of the genitourinary system. It is a Pre-mRNA-splicing regulat protein. It is a regulatory subunit of the WMM N6-methyltransferase complex which plays a role in the efficiency o mRNA splicing and processing and mRNA stability. It is required for accumulation of METTL3 and METTL14 to nucle speckle. MW of WTAP is 45-50 kDa due to phosphorylation.			
Notable Publications	Author	Pubmed ID	Journal	Application
	Fanpeng Zhao	34593014	Mol Neurodegener	WB
	Jing Wang	32998774	J Ovarian Res	WB
	Qian Luo	36175708	Med Oncol	IHC,WB


For technical support and original validation data for this product please contact: E: Proteintech-CN@ptglab.com T: 4006900926 W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


## Selected Validation Data




Immunohistochemical analysis of paraffinembedded human colon tissue slide using 60188-1-Ig (WTAP antibody) at dilution of 1:1000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).




IP result of anti-WTAP (IP:60188-1-Ig, 4ug; Detection:10200-1-AP 1:8000) with HeLa cells lysate 2000 ug.



Immunofluorescent analysis of (4% PFA ) fixed HeLa cells using 60188-1-1g(WTAP antibody) at dilution of 1:50 and Alexa Fluor 488-conjugated AffiniPure Goat Anti-Mouse IgG(H+L).



Various lysates were subjected to SDS PAGE followed by western blot with 60188-1-lg (WTAP antibody) at dilution of 1:20000 incubated at room temperature for 1.5 hours.



Various lysates were subjected to SDS PAGE followed by western blot with 60188-1-Ig (WTAP antibody) at dilution of 1:20000 incubated at room temperature for 1.5 hours. The membrane was stripped and reblotted with HRP-conjugated GAPDH Monoclonal antibody (HRP-60004) as loading control.