For Research Use Only

MYH9 Monoclonal antibody

Catalog Number:60233-1-lg Featured Product

7 Publications

BC011915

4627

P35579

227 kDa Observed MW: 230 kDa, 160 kDa

GeneID (NCBI):

UNIPROT ID:

Full Name:

Calculated MW:

GenBank Accession Number:

myosin, heavy chain 9, non-muscle



Basic Information

Applications

Catalog Number: 60233-1-lg Size:

1000 µg/ml Source: Mouse Isotype: lgG1

Immunogen Catalog Number:

AG1612

Tested Applications: IF/ICC, WB, ELISA

Cited Applications: CoIP, IF, IHC, WB Species Specificity: human, mouse, pig **Cited Species:**

human, mouse

Positive Controls:

WB: mouse kidney tissue, human kidney tissue, pig spleen tissue, U2OS cells, mouse colon tissue, pig pancreas tissue, A431 cells, human placenta tissue, pig

Purification Method:

Protein A purification

Recommended Dilutions:

WB 1:2000-1:10000 IF 1:50-1:500

CloneNo.:

5D9D2

kidney tissue

IF: HepG2 cells, U2OS cells

Background Information

MYH9 (Myosin-9) appears to play a role in cytokinesis, cell shape, and specialized functions such as secretion and capping. MYH9 is a hexameric protein that consists of two heavy chains (~200 kDa) and two pairs of non-identical light chains (15-22 kDa), a pair of essential or alkali light chains, and a pair of regulatory light chains (PMID: 12800156).

Notable Publications

Author	Pubmed ID	Journal	Application
Yiyi Liu	31594754	EBioMedicine	WB,IHC,IF
Shunfeng Hu	34743203	Cell Death Differ	WB,CoIP,IF
Gao-Shan Yang	32685020	Theranostics	IF


Storage

Store at -20°C. Stable for one year after shipment.


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


Selected Validation Data




mouse kidney tissue were subjected to SDS PAGE followed by western blot with 60233-1-1g (MYH9 antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours.



Immunofluorescent analysis of (-20°C Ethanol) fixed HepG2 cells using 60233-1-lg (MYH9 antibody) at dilution of 1:100 and CoraLite488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L).



Various lysates were subjected to SDS PAGE followed by western blot with 60233-1-1g (MYH9 antibody) at dilution of 1:20000 incubated at room temperature for 1.5 hours.



mouse colon tissue were subjected to SDS PAGE followed by western blot with 60233-1-1g (MYH9 antibody) at dilution of 1:20000 incubated at room temperature for 1.5 hours.