For Research Use Only

GOT1 Monoclonal antibody

Catalog Number:60317-1-lg 2 Publications





Basic Information	Catalog Number: 60317-1-lg	GenBank Accession Number: BC000498		Purification Method: Protein G purification	
	Size: GeneID (NCBI): 780 μg/ml 2805			CloneNo.: 5A12E10	
	Source: Mouse	UNIPROT ID: P17174		Recommended Dilutions: WB 1:2000-1:10000	
	Isotype: IgG1 Immunogen Catalog Number:	soluble (aspartate	glutamic-oxaloacetic transaminase 1, ^F 1:200-1:800 soluble (aspartate aminotransferase		
	AG7012	1) Calculated MW: 46 kDa Observed MW: 40-46 kDa			
Applications	Tested Applications:		Positive Controls:		
	FC, IF/ICC, IHC, WB, ELISA Cited Applications: IF, IP, WB			cells, rat brain tissue, rat heart tissue, 1 tissue, K-562 cells, chicken brain, chicker	
	Species Specificity: human, mouse, rat, chicken	IHC : human live		iver cancer tissue,	
	Cited Species: human			ls,	
	Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0				
Background Information	5	and L-glutamate and pla lar levels of reactive o	ays a key role in ca vygen species (ROS	rbon and nitrogen metabolism. GOT1 car) through NADPH synthesis and enhances	
Notable Publications	Author	Pubmed ID J	ournal	Application	
	Wenzhu Li	28544376 J	Cell Mol Med	WB	
	Chuanzhen Yang	37291265 N	lat Cell Biol	WB,IF,IP	
Storage	Storage: Store at -20°C. Stable for one yea Storage Buffer: PBS with 0.02% sodium azide ar Aliquoting is unnecessary for -20	d 50% glycerol pH 7.3.			

For technical support and original validation data for this product please contact: E: Proteintech-CN@ptglab.com T: 4006900926 W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data






Various lysates were subjected to SDS PAGE followed by western blot with 60317-1-1g (GOT1 antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours.


Immunohistochemical analysis of paraffinembedded human liver cancer tissue slide using 60317-1-1g (GOT1 Antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded human liver cancer tissue slide using 60317-1-lg (GOT1 Antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



1X10^6 HepG2 cells were stained with 0.20ug GOT1 antibody (60317-1-1g, red) and control antibody (blue). Fixed with 90% MeOH.



Immunofluorescent analysis of (-20°C Ethanol) fixed HepG2 cells using GOT1 antibody (60317-1-Ig, Clone: 5A12E10) at dilution of 1:400 and CoraLite®488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L).