For Research Use Only

P-selectin / CD62P Monoclonal antibody



Catalog Number:60322-1-lg


21 Publications


Basic Information	Catalog Number: 60322-1-lg	GenBank Accession Number: BC028067	Purification Method: Protein A purification	
	Size:	GenelD (NCBI):	CloneNo.:	
	1295 µg/ml	6403	3G2C7	
	Source: Mouse	UNIPROT ID: P16109	Recommended Dilutions: WB 1:2000-1:10000	
	Isotype: IgG2b Immunogen Catalog Number: AG4124	Full Name: IP 0.5-4.0 ug for 1.0-3.0 mg of total selectin P (granule membrane protein protein lysate 140kDa, antigen CD62) Calculated MW: 830 aa, 91 kDa		
				Observed MW: 140 kDa, 105-110 kDa
		Applications	Tested Applications:	Positive Controls:
IP, WB, ELISA	WB : human peripheral blood platelets, Jurkat cells			
Cited Applications: WB, IF, FC, IHC	IP : K-562 cells,			
Species Specificity: human				
Cited Species: human, rat, mouse				
	P-selectin, also known as GMP140 or CD62P, is a transmembrane glycoprotein that mediates the interaction of activated endothelial cells or platelets with leukocytes. It is an adhesion molecule involved in the pathogenesis of inflammation, thrombosis, and oncogenesis. P-selectin is stored in the alpha-granules of platelets and Weibel-Palade bodies of endothelial cells. Upon cell activation by agonists, P-selectin is transported rapidly to the cell surface.			
Background Information	activated endothelial cells or pla inflammation, thrombosis, and or Palade bodies of endothelial cell	cogenesis. P-selectin is stored in the	alpha-granules of platelets and Weibel-	
	activated endothelial cells or pla inflammation, thrombosis, and or Palade bodies of endothelial cell surface.	cogenesis. P-selectin is stored in the a s. Upon cell activation by agonists, P-s	alpha-granules of platelets and Weibel- selectin is transported rapidly to the cell	
	activated endothelial cells or pla inflammation, thrombosis, and or Palade bodies of endothelial cell surface. Author	cogenesis. P-selectin is stored in the association by agonists, P-selection by agonists, P-selec	alpha-granules of platelets and Weibel- selectin is transported rapidly to the cell Application	
	activated endothelial cells or pla inflammation, thrombosis, and or Palade bodies of endothelial cells surface. Author Jinwei Pang	Pubmed ID Journal 34566870 Front Neurol	alpha-granules of platelets and Weibel- selectin is transported rapidly to the cell Application WB,IF	
Background Information Notable Publications	activated endothelial cells or pla inflammation, thrombosis, and or Palade bodies of endothelial cells surface. Author Jinwei Pang LinXi Cheng	cogenesis. P-selectin is stored in the association by agonists, P-selection by agonists, P-selec	alpha-granules of platelets and Weibel- selectin is transported rapidly to the cell Application WB,IF FC	

For technical support and original validation data for this product please contact:T: 4006900926E: Proteintech-CN@ptglab.comW: ptgcn.com


This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data





human peripheral blood platelets were subjected to SDS PAGE followed by western blot with 60322-1-Ig (P-selectin / CD62P antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours.



IP result of anti-P-selectin / CD62P (IP:60322-1-Ig, 5ug; Detection:60322-1-Ig 1:600) with K-562 cells lysate 1760ug.

Untreated and PNGase F-treated lysates of human platelets were subjected to SDS PAGE followed by western blot with 60322-1-1g (P-selectin / CD62P antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours. PNGase F was obtained from Atagenix (cat.NO. ata808).