For Research Use Only

## Anti-Human CD3 (Hit3a)

Catalog Number:65112-1-lg





Basic Information	Catalog Number: 65112-1-lg	GenBank Accession Number: BC049847	Purification Method: Affinity purification
	Size: 500ug, 0.5 mg/ml	GenelD (NCBI): 916	CloneNo.: Hit3a
	Source: Mouse	ENSEMBL Gene ID: ENSG00000198851	
	<mark>Isotype:</mark> IgG2a, kappa	UNIPROT ID: P07766	
		Full Name: CD3e molecule, epsilon (CD3-TCR complex)	
		Calculated MW: 207 aa, 23 kDa	
Applications	Tested Applications: FC		
	Species Specificity: Human		
Background Information	CD3 is a multimeric protein associated with the T-cell receptor (TCR) to form a complex involved in antigen recognition and signal transduction (PMID: 15885124). CD3 is composed of CD3 $\gamma$ , $\delta$ , $\varepsilon$ , and $\zeta$ chains (PMID: 1826255). It is expressed by thymocytes in a developmentally regulated manner, T cells, and some NK cells (PMID: 3289580). The TCR recognizes antigens bound to major histocompatibility complex (MHC) molecules. TCR-mediated peptide-MHC recognition is transmitted to the CD3 complex, leading to the intracellular signal transduction (PMID: 11985657).		
Storage	Storage: Store at 2-8°C. Stable for one Storage Buffer: PBS with 0.09% sodium azid		

For technical support and original validation data for this product please contact:T: 4006900926E: Proteintech-CN@ptglab.comW: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

## **Selected Validation Data**





1X10<sup>^</sup>6 human PBMCs were surface stained with 0.2 ug Anti-Human CD3 (65112-1-1g, Clone:Hit3a) or Mouse IgG2a Isotype Control and CoraLite@594-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L) at dilution 1:500, then CL750 Anti-Human CD45. Cells were not fixed. Lymphocytes were gated.

1X10^6 human PBMCs were surface stained with 0.2 ug Anti-Human CD3 (65112-1-1g, Clone:Hit3a) or Mouse IgG2a Isotype Control and CoraLite@594-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L) at dilution 1:500. Cells were not fixed. Lymphocytes were gated.