For Research Use Only

Anti-Human CD11c (BU15)

Catalog Number:65196-1-lg





Basic Information	Catalog Number: 65196-1-lg	GenBank Accession Number: BC038237	Purification Method: Protein G purification
	Size: 100ug, 500 µg/ml	GenelD (NCBI): 3687	CloneNo.: BU15
	Source: Mouse	ENSEMBL Gene ID: ENSG00000140678	
	lsotype: IgG1, kappa	UNIPROT ID: P20702	
		Full Name: integrin, alpha X (complement component 3 receptor 4 subunit)	
		Calculated MW: 1169 aa, 129 kDa	
Applications	Tested Applications: FC		
	Species Specificity: Human		
Background Information	Integrins are cell adhesion receptors that are heterodimers composed of non-covalently associated α and β subunits (PMID: 9779984). CD11c, also known as integrin α X, is a type I transmembrane glycoprotein present on a variety of cells, including monocytes/macrophages, granulocytes, a subset of B cells, NK cells and dendritic cells (PMID: 2897326; 1680915; 1694698; 17389580). As a result of its high level of expression on most dendritic cells, CD11c is typically considered to be a marker of conventional dendritic cells (PMID: 27119555). CD11c forms an α / β heterodimer with CD18 (integrin β 2). CD11c/CD18 acts as a receptor for fibrinogen and is important in monocyte adhesion and chemotaxis (PMID: 1671533).		
Storage	Storage: Store at 2-8°C. Stable for one Storage Buffer: PBS with 0.09% sodium azid		

For technical support and original validation data for this product please contact:T: 4006900926E: Proteintech-CN@ptglab.comW: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data





1X10^{^6} human PBMCs were surface stained with 0.2 ug Anti-Human CD11c (65196-1-lg, Clone:BU15) and CoraLite®594-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L) at dilution 1:1000 or unstained. Cells were not fixed. Monocytes were gated.

1X10^6 human PBMCs were surface stained with 0.2 ug Anti-Human CD11c (65196-1-Ig, Clone:BU15) and CoraLite®594-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L) at dilution 1:1000 (red) or unstained. Cells were not fixed. Monocytes were gated.