For Research Use Only

CD68 Monoclonal antibody

Catalog Number: 66231-2-lg 46 Publications



Basic Information

Catalog Number:

GenBank Accession Number: BC015557

Purification Method:

66231-2-lg

Protein A purification

Size: Source: GeneID (NCBI):

CloneNo.:

1000 µg/ml

UNIPROT ID:

3A9A7

Mouse Isotype: P34810 Full Name: Recommended Dilutions: IHC 1:1000-1:6000

lgG1

CD68 molecule

Calculated MW:

Immunogen Catalog Number:

IF 1:3000-1:12000

AG22815

37 kDa

Applications

Tested Applications:

IF-P, IHC, ELISA

Cited Applications:

WB,IHC,IF,FC

Species Specificity:

human

Cited Species: human, pig

Positive Controls:

IHC: human tonsillitis tissue, human lung cancer tissue, human liver cancer tissue, human urothelial carcinoma tissue, human appendicitis tissue, human

colon cancer tissue

IF: human tonsillitis tissue,

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate

buffer pH 6.0

Background Information

CD68 is a type I transmembrane glycoprotein that is highly expressed by human monocytes and tissue macrophages. It belongs to the lysosomal/endosomal-associated membrane glycoprotein (LAMP) family and primarily localizes to lysosomes and endosomes with a smaller fraction circulating to the cell surface. CD68 is also a member of the scavenger receptor family. It may play a role in phagocytic activities of tissue macrophages.

Notable Publications

| Author | Pubmed ID | Journal | Application |
|---------------|-----------|------------------|-------------|
| Yu Huang | 34713723 | J Am Heart Assoc | |
| Yourong Jiang | 36389736 | Front Immunol | IF |
| Xinyi Zhang | 36499018 | Int J Mol Sci | IF |


Storage

Store at -20°C. Stable for one year after shipment.


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


Selected Validation Data




Immunohistochemical analysis of paraffinembedded human tonsillitis tissue slide using 66231-2-Ig (CD68 antibody) at dilution of 1:2000 (under 10x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded human tonsillitis tissue slide using 66231-2-Ig (CD68 antibody) at dilution of 1:2000 (under 40x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (4% PFA) fixed human tonsillitis tissue using CD68 antibody (66231-2-Ig, Clone: 3A9A7) at dilution of 1:6000 and Coralite®488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L).



Immunofluorescent analysis of (4% PFA) fixed human tonsillitis tissue using CD68 antibody (66231-2-lg, Clone: 3A9A7) at dilution of 1:6000 and CoraLite®488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L).