For Research Use Only

CD68 Monoclonal antibody

Catalog Number:66231-2-lg 46 Publications




Basic Information	Catalog Number: 66231-2-lg	GenBank Accession Num BC015557	ber: Purification Method: Protein A purification				
	Size:	GeneID (NCBI):	CloneNo.:				
	1000 µg/ml	968	3A9A7				
	Source:	UNIPROT ID:	Recommended Dilutions:				
	Mouse	P34810	IHC 1:1000-1:6000				
	lsotype: lgG1	Full Name: CD68 molecule	IF 1:3000-1:12000				
	Immunogen Catalog Number: AG22815	Calculated MW: 37 kDa					
Applications	Tested Applications: IF-P, IHC, ELISA						
			IC : human tonsillitis tissue, human lung cancer ssue, human liver cancer tissue, human urothelial				
	IF, FC, IHC		carcinoma tissue, human appendicitis tissue, human colon cancer tissue				
	Species Specificity:						
	human	IF	: human tonsillitis tissue,				
	Cited Species: human, pig Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0						
					with citrate buffer pH 6.	o 0	
				Background Information	CD68 is a type I transmembrane g macrophages. It belongs to the ly primarily localizes to lysosomes	O glycoprotein that is highly exp sosomal/endosomal-associat and endosomes with a smalle	ressed by human monocytes and tissue ed membrane glycoprotein (LAMP) family and r fraction circulating to the cell surface. CD68 is als n phagocytic activities of tissue macrophages.
	CD68 is a type I transmembrane g macrophages. It belongs to the ly primarily localizes to lysosomes	O glycoprotein that is highly exp sosomal/endosomal-associat and endosomes with a smalle	ed membrane glycoprotein (LAMP) family and r fraction circulating to the cell surface. CD68 is als				
	With citrate buffer pH 6. CD68 is a type I transmembrane g macrophages. It belongs to the ly primarily localizes to lysosomes a member of the scavenger recep	O glycoprotein that is highly exp sosomal/endosomal-associat and endosomes with a smalle tor family. It may play a role i	ed membrane glycoprotein (LAMP) family and r fraction circulating to the cell surface. CD68 is als n phagocytic activities of tissue macrophages. Application				
Background Information Notable Publications	With citrate buffer pH 6. CD68 is a type I transmembrane g macrophages. It belongs to the ly primarily localizes to lysosomes a member of the scavenger recep	Q Rycoprotein that is highly exp sosomal/endosomal-associat and endosomes with a smalle tor family. It may play a role i Pubmed ID Journal	ed membrane glycoprotein (LAMP) family and r fraction circulating to the cell surface. CD68 is als n phagocytic activities of tissue macrophages. Application				
	With citrate buffer pH 6. CD68 is a type I transmembrane g macrophages. It belongs to the ly primarily localizes to lysosomes a member of the scavenger recep Author Yu Huang	Q glycoprotein that is highly exposored sosomal/endosomal-associat and endosomes with a smalle tor family. It may play a role i Pubmed ID Journal 34713723 J Am Heat	ed membrane glycoprotein (LAMP) family and r fraction circulating to the cell surface. CD68 is als n phagocytic activities of tissue macrophages. Application art Assoc munol IF				
	with citrate buffer pH 6. CD68 is a type I transmembrane g macrophages. It belongs to the ly primarily localizes to lysosomes a member of the scavenger recep Author Yu Huang Yourong Jiang	Q glycoprotein that is highly exposored leadsomal-associat and endosomes with a smaller tor family. It may play a role is Pubmed ID Journal 34713723 J Am Heir 36389736 Front Im 36499018 Int J Mol	ed membrane glycoprotein (LAMP) family and r fraction circulating to the cell surface. CD68 is als n phagocytic activities of tissue macrophages. Application art Assoc nunol IF				


For technical support and original validation data for this product please contact: E: Proteintech-CN@ptglab.com T: 4006900926 W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




Immunohistochemical analysis of paraffinembedded human tonsillitis tissue slide using 66231-2-Ig (CD68 antibody) at dilution of 1:2000 (under 10x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded human tonsillitis tissue slide using 66231-2-Ig (CD68 antibody) at dilution of 1:2000 (under 40x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (4% PFA) fixed human tonsillitis tissue using CD68 antibody (66231-2-Ig, Clone: 3A9A7) at dilution of 1:6000 and Coralite®488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L).



Immunofluorescent analysis of (4% PFA) fixed human tonsillitis tissue using CD68 antibody (66231-2-1g, Clone: 3A9A7) at dilution of 1:6000 and CoraLite®488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L).