

PD-L1/CD274 Monoclonal antibody

Catalog Number: 66248-1-Ig

Featured Product

245 Publications

Basic Information

Catalog Number:

66248-1-Ig

Size:

2000 µg/ml

Source:

Mouse

Isotype:

IgG1

Immunogen Catalog Number:

AG12443

GenBank Accession Number:

BC074984

GeneID (NCBI):

29126

UNIPROT ID:

Q9NZQ7

Full Name:

CD274 molecule

Calculated MW:

290 aa, 33 kDa

Observed MW:

45-50 kDa

Purification Method:

Protein A purification

CloneNo.:

2B11D11

Recommended Dilutions:

WB 1:2000-1:10000

IHC 1:5000-1:20000

IF 1:50-1:500

Applications

Tested Applications:

FC, IF/ICC, IHC, WB, ELISA

Cited Applications:

CoIP, FC, IF, IHC, IP, WB

Species Specificity:

human, mouse, pig, rat

Cited Species:

human, rat, mouse

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB: A375 cells, human placenta tissue, pig lung tissue, human skeletal muscle tissue, HepG2 cells, THP-1 cells, RAW 264.7 cells, A549 cells, K-562 cells, HSC-T6 cells

IHC: human tonsillitis tissue, human heart tissue, human lung cancer tissue, human placenta tissue, mouse heart tissue

IF: HeLa cells,

Background Information

Programmed cell death ligand 1 (PD-L1, CD274, or B7-H1), is the first member of B7 family to be discovered. B7 family molecules are type I transmembrane proteins belonging to the immunoglobulin superfamily. In concert with their CD28 family receptors, the B7s are key regulators of the adaptive immune response. PD-L1 is suggested as a negative regulator of T and B cell, and plays important role in mediating tolerance of lymphocytes to self-antigens. It is also involved in the costimulatory signal, essential for T-cell proliferation and production of IL10 and IFNG, in an IL2-dependent and a PD-1-independent manner. PD-L1 is a 290 aa transmembrane protein with a calculated molecular weight of 33 kDa, it is predicted to be 27-30 kDa after signal peptide cleavage (PMID: 25609200; 17076679). The apparent molecular weight has also been reported as 45-70 kDa, major glycosylated form of 45-50 kDa and multiple post-translational modifications form of 65-70 kDa (PMID: 18760278; 16493058).

Notable Publications

Author	Pubmed ID	Journal	Application
Jiacheng Huang	34650926	Front Oncol	IHC
Youqiong Ye	32988398	Genome Med	WB
Hao Zhang	36136350	Brief Bioinform	IHC

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


For technical support and original validation data for this product please contact:

T: 4006900926


E: Proteintech-CN@ptglab.comW: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




human placenta tissue were subjected to SDS PAGE followed by western blot with 66248-1-Ig (PD-L1/CD274 antibody) at dilution of 1:4000 incubated at room temperature for 1.5 hours.




Untreated and IFN gamma treated A375 cells and A549 cells were subjected to SDS PAGE followed by western blot with 66248-1-Ig (PD-L1/CD274 antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours.




Immunohistochemical analysis of paraffin-embedded human tonsillitis tissue slide using 66248-1-Ig (PD-L1/CD274 antibody) at dilution of 1:10000 (under 10x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffin-embedded human tonsillitis tissue slide using 66248-1-Ig (PD-L1/CD274 antibody) at dilution of 1:10000 (under 4x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (10% Formaldehyde) fixed HeLa cells using 66248-1-Ig(PD-L1/CD274 antibody) at dilution of 1:300 and Alexa Fluor 488-conjugated AffiniPure Goat Anti-Mouse IgG(H+L).



1X10⁶ MDA-MB-231 cells were stained with 0.2 ug Anti-Human PD-L1/CD274 (66248-1-Ig, Clone: 2B11D11) and CoraLite®488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L) at dilution 1:1000 (green), or stained with 0.2 ug mouse IgG1 isotype control antibody and CoraLite®488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L) at dilution 1:1000 (black). Cells were fixed with 90% MeOH.