### For Research Use Only

# PEX5 Monoclonal antibody

Catalog Number:66309-1-lg

**Featured Product** 

1 Publications

BC010621

5830

P50542

GeneID (NCBI):

**UNIPROT ID:** 

Full Name:



**Basic Information** 

Catalog Number: 66309-1-lg

Size: 1500 µg/ml Source: Mouse Isotype:

lgG1 Immunogen Catalog Number:

AG3268

Calculated MW: 631 aa, 70 kDa

GenBank Accession Number:

peroxisomal biogenesis factor 5

Observed MW: 70-80 kDa

**Applications** 

**Tested Applications:** IHC, WB, ELISA **Cited Applications:** 

WB

Species Specificity: human, pig, rat **Cited Species:** human

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (\*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB: pig brain tissue, HEK-293 cells, fetal human brain

**Purification Method:** 

Protein G purification

Recommended Dilutions:

WB 1:500-1:2000 IHC 1:50-1:500

CloneNo.:

1C2C3

IHC: human liver tissue, human kidney tissue

## **Background Information**

The peroxisomal targeting signal type1 (PTS1) receptor, PEX5, is one member of peroxins (PEXs) which are proteins required for peroxisome assembly. PEX5 and PEX7 function as receptors that recognize PTS1- and PTS2- containing proteins, respectively, and PEX5 binds PTS1 through its C-terminal 40-kDa tetratricopeptide repeat domain. It is a predominantly cytoplasmic, partly peroxisomal protein that appears to shuttle between these compartments as it mediates the import of PTS1-containing proteins. PEX5 has been reported to interact with PEX10, PEX12, PEX13, and PEX14. Defects in PEX5 are a cause of Zellweger syndrome (ZWS), which is a lethal peroxisome biogenesis disorder.

#### **Notable Publications**

Léa P Wilhelm 36215693 EMBO J WB	

Storage


Storage:

Store at -20°C. Stable for one year after shipment.


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

## **Selected Validation Data**




WB result of PEX5 antibody (66309-1-lg; 1:1000; incubated at room temperature for 1.5 hours) with sh-Control and sh-PEX5 transfected HEK-293 cells.



pig brain tissue were subjected to SDS PAGE followed by western blot with 66309-1-1g (PEXS Antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded human liver tissue slide using 66309-1-Ig (PEX5 Antibody) at dilution of 1:200 (under 10x lens).



Immunohistochemical analysis of paraffinembedded human liver tissue slide using 66309-1-Ig (PEX5 Antibody) at dilution of 1:200 (under 40x lens).