## For Research Use Only

## GATA3 Monoclonal antibody, PBS Only



**Purification Method:** 

Protein A purification

CloneNo.:

1B11A8

Catalog Number: 66400-1-PBS

**Basic Information** 

Catalog Number: 66400-1-PBS

Size:

GenBank Accession Number:

BC003070

GeneID (NCBI):

2625

1 mg/ml Source: **UNIPROT ID:** 

Mouse P23771 Full Name: Isotype:

lgG2a GATA binding protein 3

Calculated MW: Immunogen Catalog Number: AG0174

443 aa, 48 kDa Observed MW:

50 kDa

**Applications** 

**Tested Applications:** 

WB,Indirect ELISA,IHC,IF

Species Specificity:

human, mouse

**Background Information** 

GATA3 is a transcriptional factor that binds to the enhancer elements of all 4T-cell antigen receptor genes. GATA3 is highly expressed in naive, freshly activated cells and Th2 lineage cells and thought to be necessary and sufficient for Th2 cytokine gene expression. GATA3 and TBET interfer with each other to regulate gene expression.

Storage

Storage:


Store at -80°C.

Storage Buffer:


## Selected Validation Data




MCF-7 cells were subjected to SDS PAGE followed by western blot with 66400-1-Ig (GATA3 Antibody) at dilution of 1:3000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 66400-1-PBS in a different storage buffer formulation.



Immunohistochemical analysis of paraffinembedded human breast cancer tissue slide using 66400-1-lg (GATA3 Antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 66400-1-PBS in a different storage buffer formulation.



Immunohistochemical analysis of paraffinembedded human colon cancer tissue slide using 66400-1-Ig (GATA3 Antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 66400-1-PBS in a different storage buffer formulation.



Immunofluorescent analysis of (4% PFA) fixed MCF-7 cells using GATA3 antibody (66400-1-1g, Clone: 1811A8) at dilution of 1:800 and CoraLite® 488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L). This data was developed using the same antibody clone with 66400-1-PBS in a different storage buffer formulation.