

HVCN1 Monoclonal antibody

Catalog Number: 66449-1-Ig

Basic Information

Catalog Number: 66449-1-Ig	GenBank Accession Number: BC032672	Purification Method: Protein G purification
Size: 1800 µg/ml	GeneID (NCBI): 84329	CloneNo.: 1E4C4
Source: Mouse	UNIPROT ID: Q96D96	Recommended Dilutions: WB 1:2000-1:16000 IHC 1:50-1:500 IF 1:200-1:800
Isotype: IgG1	Full Name: hydrogen voltage-gated channel 1	
Immunogen Catalog Number: AG5350	Calculated MW: 273 aa, 32 kDa Observed MW: 28-35 kDa, 40 kDa, 60 kDa	

Applications

Tested Applications: IF-P, IHC, WB, ELISA	Positive Controls:
Species Specificity: human	WB : Raji cells, HeLa cells
Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0	IHC : human tonsillitis tissue,
	IF : human tonsillitis tissue,

Background Information

HVCN1, also named as VSOP and HV1, Belongs to the hydrogen channel family. HVCN1 mediates the voltage-dependent proton permeability of excitable membranes. It forms a proton-selective channel through which protons may pass in accordance with their electrochemical gradient. Proton efflux, HVCN1 is accompanied by membrane depolarization, facilitates acute production of reactive oxygen species in phagocytosis. HVCN1, the voltage-sensitive proton channel, is present in human sperm and is an important regulator of the functional maturation of sperm. HVCN1 has four isoforms with MW 28-32 kDa or 40 kDa (modification). It has a dimer form with MW 60 kDa.

Storage

Storage:
Store at -20°C. Stable for one year after shipment.
Storage Buffer:
PBS with 0.02% sodium azide and 50% glycerol pH 7.3.
 Aliquoting is unnecessary for -20°C storage


For technical support and original validation data for this product please contact:

T: 4006900926


E: Proteintech-CN@ptglab.comW: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




Raji cells were subjected to SDS PAGE followed by western blot with 66449-1-Ig (HVCN1 antibody) at dilution of 1:8000 incubated at room temperature for 1.5 hours.




Immunofluorescent analysis of (4% PFA) fixed human tonsillitis tissue using HVCN1 antibody (66449-1-Ig, Clone: 1E4C4) at dilution of 1:400 and CoraLite®488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L).



Immunofluorescent analysis of (4% PFA) fixed human tonsillitis tissue using HVCN1 antibody (66449-1-Ig, Clone: 1E4C4) at dilution of 1:400 and CoraLite®488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L).



Immunohistochemical analysis of paraffin-embedded human tonsillitis tissue slide using 66449-1-Ig (HVCN1 Antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffin-embedded human tonsillitis tissue slide using 66449-1-Ig (HVCN1 Antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).