For Research Use Only

EGFR Monoclonal antibody, PBS Only

Catalog Number: 66455-1-PBS

Featured Product



Basic Information

Catalog Number:

BC094761

Purification Method:

66455-1-PBS

GeneID (NCBI):

Protein G purification

Size: 1 mg/ml

1956

CloneNo.:

Source:

UNIPROT ID:

P00533

2A2H10

Mouse Isotype:

Full Name:

lgG1 Immunogen Catalog Number:

epidermal growth factor receptor (erythroblastic leukemia viral (v-erb-

b) oncogene homolog, avian)

GenBank Accession Number:

AG24947

Calculated MW:

1210 aa, 134 kDa Observed MW:

145-165 kDa

Applications

Tested Applications:

WB, FC, IHC, Indirect ELISA

Species Specificity:

human

Background Information


EGFR, also named as ERBB1, is a cell-surface receptor for members of the epidermal growth factor family (EGFfamily) of extracellular protein ligands. Binding of the protein to a ligand induces receptor dimerization and tyrosine autophosphorylation and leads to cell proliferation. The gene resides on chromosome 7p12, encoding a 170 kDa membrane-associated glycoprotein. Recent studies have shown EGFR plays a critical role in cancer development and progression, including cell proliferation, apoptosis, angiogenesis, and metastatic spread. Mutations in this gene are associated with lung cancer.

Storage


Storage: Store at -80°C.

Storage Buffer: PBS Only

Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 66455-1-lg (EGFR antibody) at dilution of 1:20000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 66455-1-PBS in a different storage buffer formulation.




WB result of EGFR antibody (66455-1-Ig; 1:20000; incubated at room temperature for 1.5 hours) with sh-Control and sh-EGFR transfected HepG2 cells. This data was developed using the same antibody clone with 66455-1-PBS in a different storage buffer formulation.



Immunohistochemical analysis of paraffinembedded human tonsillitis tissue slide using 66455-1-lg (EGFR antibody) at dilution of 1:2000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 66455-1-PBS in a different storage buffer formulation.



Immunohistochemical analysis of paraffinembedded human tonsillitis tissue slide using 66455-1-lg (EGFR antibody) at dilution of 1:2000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 66455-1-PBS in a different storage buffer formulation.



1X10^6 A431 cells were intracellularly stained with 0.4 ug Anti-Human EGFR (66455-1-lg, Clone:2A2H10) and Coralite®488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L) at dilution 1:1000 (red), or 0.4 ug Mouse IgG1 Isotype Control (MOPC-21) (65124-1-lg, Clone: MOPC-21) (blue). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C). This data was developed using the same antibody clone with 66455-1-PBS in a

