

For Research Use Only

MDM2 Monoclonal antibody

Catalog Number: 66511-1-Ig **21 Publications**



Basic Information

Catalog Number: 66511-1-Ig	GenBank Accession Number: NM_002392	Purification Method: Protein A purification
Size: 2000 µg/ml	GeneID (NCBI): 4193	CloneNo.: 2A6C9
Source: Mouse	UNIPROT ID: Q00987	Recommended Dilutions: WB 1:1000-1:6000 IHC 1:150-1:600 IF 1:400-1:1600
Isotype: IgG2b	Full Name: Mdm2 p53 binding protein homolog (mouse)	
	Calculated MW: 55 kDa	
	Observed MW: 40-55 kDa	

Applications

Tested Applications:
IF/ICC, IHC, WB, ELISA

Cited Applications:
CoIP, IF, IHC, IP, WB

Species Specificity:
Human, mouse, rat, rabbit

Cited Species:
human, rat, mouse

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB: MCF-7 cells, A549 cells, mouse brain tissue, rabbit brain tissue, rat brain tissue, U-251 cells

IHC: human breast cancer tissue, human liver cancer tissue

IF: HeLa cells,

Background Information

MDM2, also known as HDM2, is a protein of the E3 ubiquitin-ligase family. MDM2 is recognized as a proto-oncogene and is over-expressed in a wide range of human malignancies including soft tissue carcinomas and breast cancer. MDM2 can bind to the p53 tumor suppressor protein with high affinity, and negatively regulates p53 by mediating its ubiquitination and proteosomal degradation (PMID: 17000718). MDM2 has some isoforms with calculated molecular mass of 25-55kD, and can be detected as 90-97kD. In addition to full length MDM2, two lower bands (85kD and 75kD) are often observed which are likely to be produced by translation initiation at methionine codons 50 and 60. Another isoform of MDM2 is often observed as 57-60kD in cell lines which is indeed a caspase cleavage product of MDM2 at residue 361 (PMID: 19565011, 9840926, 7689721).

Notable Publications

Author	Pubmed ID	Journal	Application
Yanting Zhu	34552711	Pulm Circ	WB
Suping Li	34687132	J Cell Mol Med	IF
Jingwen Tan	36208777	Chem Biol Interact	WB


Storage

Storage:
Store at -20°C. Stable for one year after shipment.
Storage Buffer:
PBS with 0.02% sodium azide and 50% glycerol pH 7.3.
Aliquoting is unnecessary for -20°C storage


For technical support and original validation data for this product please contact:
T: 4006900926 E: Proteintech-CN@ptglab.com W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




MCF-7 cells were subjected to SDS PAGE followed by western blot with 66511-1-Ig (MDM2 antibody) at dilution of 1:3000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffin-embedded human breast cancer tissue slide using 66511-1-Ig (MDM2 antibody) at dilution of 1:300 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffin-embedded human breast cancer tissue slide using 66511-1-Ig (MDM2 antibody) at dilution of 1:300 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (4% PFA) fixed HeLa cells using MDM2 antibody (66511-1-Ig, Clone: 2A6C9) at dilution of 1:800 and CoraLite@488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L).