For Research Use Only

LATS1 Monoclonal antibody

Catalog Number:66569-1-lg Featured Product



Basic Information

Catalog Number: GenBank Accession Number: 66569-1-lg BC015665 GeneID (NCBI): Size: 700 µg/ml 9113 **UNIPROT ID:** Source: Mouse 095835

Full Name: Isotype: lgG1 LATS, large tumor suppressor,

homolog 1 (Drosophila) Immunogen Catalog Number: AG11140

Calculated MW:

15 kDa, 127 kDa Observed MW: 140-150 kDa, 120 kDa **Purification Method:**

Protein G purification CloneNo.:

2A8F1

Recommended Dilutions: WB 1:2000-1:10000

Applications

Tested Applications: WB, ELISA

Species Specificity:

Human

Positive Controls:

WB: T-47D cells, HeLa cells, HEK-293 cells, HepG2 cells, MCF-7 cells, K-562 cells, Jurkat cells

Background Information

LATS1(Large tumor suppressor homolog 1) is also named as WARTS and belongds to the AGC Ser/Thr protein kinase family. The gene encodes a highly conserved (from fly to human) protein kinase that plays a crucial role in the $prevention of tumor formation by controlling the progression of mitosis. The expression of both long (170 \, kDa) and$ short lats1 isoforms(120 kDa) in vertebrate retinal cells raises the possibility that these lats1 proteins may act as negative key regulators of the cell cycle each of them performing a unique role (PMID:15777619). In mammalian cells, LATS1 was phosphorylated in a cell cycle-dependent manner and complexed with CDC2 in early mitosis (PMID:9988268). LATS1 also can be detected as 120 kDa and 140-150 kDa, and play a key role in the regulation of Hippo pathway (PMID: 27940445).

Storage


Storage:

Store at -20°C. Stable for one year after shipment.


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 66569-1-1g (LATS1 antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours.



WB result of LATS1 antibody (66569-1-Ig; 1:300; incubated at room temperature for 1.5 hours) with sh-Control and sh-LATS1 transfected HepG2 cells.



HepG2 cells were subjected to SDS PAGE followed by western blot with 66569-1-1g (LATS1 antibody) at dilution of 1:2000 incubated at room temperature for 1.5 hours.