For Research Use Only

ANLN Monoclonal antibody

Catalog Number:66643-1-lg Featured Product

4 Publications



Basic Information

Catalog Number: 66643-1-lg Size:

1600 µg/ml Source: Mouse Isotype:

lgG1 Immunogen Catalog Number:

AG19294

GenBank Accession Number:

BC034692 GeneID (NCBI): 54443 **UNIPROT ID:** Q9NQW6 Full Name:

anillin, actin binding protein

Calculated MW: 1124 aa, 124 kDa Observed MW: 110 kDa~120 kDa **Purification Method:** Protein A purification

CloneNo.:

1B4A8

Recommended Dilutions: WB 1:2000-1:10000

Applications

Tested Applications:

WB, ELISA

Cited Applications:

WB

Species Specificity: Human, Mouse, Rat Cited Species: human

Positive Controls:

WB: HeLa cells, MCF-7 cells, rat brain tissue, mouse brain tissue, HEK-293 cells, U-251 cells, PC-12 cells

Background Information

Anillin, an actin-binding protein, is a conserved cleavage furrow component important for cytokinesis. The human anillin protein localizes to the nucleus during interphase, the cortex following nuclear envelope breakdown, and the cleavage furrow during cytokinesis. It interactions with at least two other furrow proteins, actin and the septins through its C-terminal PH domain and N-terminal actin-binding domain. In anillin knockdown cells, the cleavage $furrow\ ingressed\ but\ failed\ to\ complete\ the\ ingression.\ An illin\ has\ been\ found\ to\ be\ overexpressed\ in\ tumors\ and$ may serve as promising marker for pancreatic carcinoma molecular diagnostics as well as a prognostic indicator for non-small cell lung cancers.

Notable Publications

Author	Pubmed ID	Journal	Application
Xiaofeng Dai	31636652	Front Genet	WB
Min Zhang	32198770	Clin Exp Pharmacol Physiol	WB
Ling Li	32275843	Cell Cycle	WB


Storage


Store at -20°C. Stable for one year after shipment.


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

Selected Validation Data







WB result of ANLN antibody (66643-1-lg; 1:3000; incubated at room temperature for 1.5 hours) with sh-Control and sh-ANLN transfected MCF-7 cells.