

OCIAD1 Monoclonal antibody

Catalog Number: 66698-1-Ig

Featured Product

1 Publications

Basic Information

Catalog Number:

66698-1-Ig

Size:

1900 µg/ml

Source:

Mouse

Isotype:

IgG2a

Immunogen Catalog Number:

AG9977

GenBank Accession Number:

BC003409

GeneID (NCBI):

54940

UNIPROT ID:

Q9NX40

Full Name:

OCIA domain containing 1

Calculated MW:

28 kDa

Observed MW:

35 kDa

Purification Method:

Protein A purification

CloneNo.:

1C10C3

Recommended Dilutions:

WB 1:1000-1:5000

IHC 1:50-1:500

IF 1:200-1:800

Applications

Tested Applications:

IF/ICC, IF-P, IHC, WB, ELISA

Cited Applications:

WB

Species Specificity:

Human, Mouse, Rat

Cited Species:

human

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB : HeLa cells, rat testis tissue, mouse testis tissue

IHC : human liver cancer tissue, human kidney tissue, human pancreas cancer tissue, human thyroid cancer tissue, mouse testis tissue

IF : human liver cancer tissue, HeLa cells

Background Information

OCIAD1 was first identified by immunoscreening of an ovarian carcinoma cDNA expression library with ascites fluid from ovarian cancer patients (PMID: 11162530). OCIAD1 has been reported as a key player in ovarian cancer cell adhesion, as well as a key player in generating ovarian cancer recurrence (PMID: 18328549; 20515946). In addition to its roles in cancer, OCIAD1 participates in maintaining stem cell potency by regulating the Jak/STAT pathway (PMID: 23972987). Several alternatively spliced forms of OCIAD1 gene have been identified. The longest form (1.4 kb) is predicted to encode for a 27.6 kDa protein of 245 amino acids. This antibody detects OCIAD1 with an apparent molecular weight of ~35 kDa as has been demonstrated by several researches (PMID: 27345969; 27345976).

Notable Publications

Author	Pubmed ID	Journal	Application
Maxence Le Vasseur	34034859	Elife	WB

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com


W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




HeLa cells were subjected to SDS PAGE followed by western blot with 66698-1-Ig (OCIAD1 antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.




Immunohistochemical analysis of paraffin-embedded human liver cancer tissue slide using 66698-1-Ig (OCIAD1 antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffin-embedded human liver cancer tissue slide using 66698-1-Ig (OCIAD1 antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (4% PFA) fixed human liver cancer tissue using OCIAD1 antibody (66698-1-Ig, Clone: 1C10C3) at dilution of 1:400 and CoraLite®488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L).



Immunofluorescent analysis of (4% PFA) fixed human liver cancer tissue using OCIAD1 antibody (66698-1-Ig, Clone: 1C10C3) at dilution of 1:400 and CoraLite®488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L).