

For Research Use Only

TRIB3 Monoclonal antibody

Catalog Number: 66702-1-Ig **2 Publications**



Basic Information

Catalog Number: 66702-1-Ig	GenBank Accession Number: BC027484	Purification Method: Protein A purification
Size: 1800 µg/ml	GeneID (NCBI): 57761	CloneNo.: 2C10C7
Source: Mouse	UNIPROT ID: Q96RU7	Recommended Dilutions: WB 1:2000-1:10000
Isotype: IgG1	Full Name: tribbles homolog 3 (Drosophila)	
Immunogen Catalog Number: AG25776	Calculated MW: 358 aa, 40 kDa	
	Observed MW: 40-45 kDa	

Applications

Tested Applications: WB, ELISA	Positive Controls: WB: HT-29 cells, MCF-7 cells, rat liver tissue, mouse hepatocytes, DU 145 cells, LNCaP cells, PC-3 cells, HUVEC cells, HepG2 cells, LO2 cells
Cited Applications: WB	
Species Specificity: Human, Mouse, Rat	
Cited Species: human, mouse	

Background Information

TRIB3 is a pseudokinase molecule that affects a number of cellular functions. TRIB3 has been reported to be highly activated in the presence of a variety of stressors, including the deprivation of neurotrophic factors, hypoxia, and ER stress. TRIB3 protein is associated with a good prognosis in human breast cancer patients, possibly due to the fact that TRIB3 is involved in hypoxia tolerance. (PMID: 21704407). TRIB3 protein predicted molecular weights is approximately 39-45 kDa. (PMID: 31191318, PMID: 24414038, PMID: 31578236).

Notable Publications

Author	Pubmed ID	Journal	Application
Hang Zhao	31118581	Drug Des Devel Ther	WB
Jean-David Morel	36752209	JCI Insight	WB

Storage

Storage:
Store at -20°C. Stable for one year after shipment.
Storage Buffer:
PBS with 0.02% sodium azide and 50% glycerol pH 7.3.
Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com


W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 66702-1-Ig (TRIB3 antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours.



Various lysates were subjected to SDS PAGE followed by western blot with 66702-1-Ig (TRIB3 antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours.