## For Research Use Only

# Cytokeratin 5 Monoclonal antibody

Catalog Number:66727-1-lg 11 Publications



**Basic Information** 

Catalog Number: 66727-1-Ig Concentration:

1000 ug/ml

Source:

AG24184

BC024292 GeneID (NCBI): 3852 UNIPROT ID: P13647

GenBank Accession Number:

Mouse P13647

Isotype: Full Name: keratin 5

Immunogen Catalog Number: Calculated MW:

Observed MW: 60 kDa

590 aa, 62 kDa

Purification Method: Protein G purification

CloneNo.: 1A1C5

Recommended Dilutions: WB 1:20000-1:100000 IHC 1:5000-1:20000 IF-P 1:200-1:800 IF/ICC 1:200-1:800

Applications

Tested Applications:

WB, IHC, IF/ICC, IF-P, ELISA

**Cited Applications:** 

WB, IHC, IF

Species Specificity: human, mouse, pig Cited Species: human, mouse, rat

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (\*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

#### **Positive Controls**

WB: HaCaT cells, SKOV-3 cells, mouse skin tissue, MCF-7 cells, A549 cells, A-253 cells, A431 cells, human saliva, pig skin tissue

IHC: human tonsillitis tissue, human breast cancer tissue, human breast hyperplasia tissue, human cervical cancer tissue, human lung cancer tissue, human oesophagus cancer tissue, human skin cancer tissue.

IF-P: human oesophagus cancer tissue, human prostate cancer tissue, human lung cancer tissue

IF/ICC: A431 cells,

# **Background Information**

Keratins are a large family of proteins that form the intermediate filament cytoskeleton of epithelial cells. Keratin expression is highly regulated, tissue specific, and varies according to cell-state. Type I keratins consist of acidic, low molecular weight proteins with MW ranging from 40 kDa (KRT19) to 64 kDa (KRT9). Type 2 keratins consist of basic or neutral, high molecular weight proteins with MW from 52 kDa (KRT8) to 67 kDa (KRT18). Keratin 5, one 58-kD type II keratin, is coexpressed with a 50-kD keratin 14 in stratified squamous epithelia. Keratin 5, one 58-kD type II keratin, is coexpressed with a 50-kD tyK14 in stratified squamous epithelia.

### **Notable Publications**

Author	Pubmed ID	Journal	Application
Kui Xiao	34714841	PLoS One	WB
Mie Naruse	34912374	Front Genet	IHC
Xin-Xing Lei	32382038	Cell Death Dis	WB

Storage

Storage

Store at -20°C. Stable for one year after shipment.

Storage Buffer

PBS with 0.02% sodium azide and 50% glycerol, pH7.3

Aliquoting is unnecessary for -20°C storage


For technical support and original validation data for this product please contact:

T: 4006900926 E: Proteintech-CN@ptglab.com


W: ptgcn.coi

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


# **Selected Validation Data**




Various lysates were subjected to SDS PAGE followed by western blot with 66727-1-lg (Cytokeratin 5 antibody) at dilution of 1:100000 incubated at room temperature for 1.5 hours.




Immunohistochemical analysis of paraffinembedded human tonsillitis tissue slide using 66727-1-1g (Cytokeratin 5 antibody) at dilution of 1:10000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).




Immunofluorescent analysis of (-20°C Methanol) fixed A451 cells using Cytokeratin 5 antibody (66727-1-1g, Clone: 1A1C5) at dilution of 1:400 and CoraLite®488-Conjugated Affini Pure Goat Anti-Mouse IgG(H+L).



Immunohistochemical analysis of paraffinembedded human tonsillitis tissue slide using 66727-1-lg (Cytokeratin 5 antibody) at dilution of 1:10000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (4% PFA) fixed human oesophagus cancer tissue using Cytokeratin 5 antibody (66727-1-1g, Clone: 1A1C5) at dilution of 1:400 and Coralite®488-Conjugated Affini Pure Goat Anti-Mouse IgG(H+L).



Immunohistochemical analysis of paraffinembedded human oesophagus cancer tissue slide using 66727-1-Ig (Cytokeratin 5 antibody) at dilution of 1:2000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).