For Research Use Only

OGT Monoclonal antibody

Catalog Number:66823-1-lg Featured Product

8 Publications



Basic Information

Catalog Number: 66823-1-lg Concentration:

1400 µg/ml

Source: Mouse Isotype: IgG2a

Immunogen Catalog Number:

AG28402

GenBank Accession Number:

BC014434 GeneID (NCBI): 8473 **UNIPROT ID:** 015294

Full Name: O-linked N-acetylglucosamine (GlcNAc) transferase (UDP-Nacetylglucosamine:polypeptide-Nacetylglucosaminyl transferase)

Calculated MW: 1046 aa, 117 kDa Observed MW: 110 kDa

Purification Method: Protein A purification

CloneNo.:

2B2A6 Recommended Dilutions: WB 1:5000-1:50000 IHC 1:250-1:1000

IF/ICC 1:200-1:800

Applications

Tested Applications: WB, IHC, IF/ICC, ELISA Cited Applications: WB, IHC, IF

Species Specificity: Human, mouse, rat, pig

Cited Species: human, mouse

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB: HeLa cells, HEK-293 cells, HSC-T6 cells, NIH/3T3 cells, pig brain tissue, rat brain tissue, mouse brain

IHC: human lung cancer tissue, human colon cancer

IF/ICC: HepG2 cells,

Background Information

 $O-linked\ N-acetylglucosamine\ transferase\ (OGT)\ catalyzes\ the\ attachment\ of\ N-acetylglucosamine\ (GlcNAc)$ monosaccharides to the hydroxyl group of serine or threonine residues of numerous nuclear and cytoplasmic proteins and may play important roles in a large number of diverse intracellular processes ranging from translational control, transcription, transcriptional repression, INS resistance and regulation of the cell cycle. It exists as a heterotrimeric complex with two 110 kDa and one 70 kDa subunits. Recent studies have shown that O-GlcNAcylation plays essential roles in cancer formation and progression. O-GlcNAcylation as well as OGT expression was found to be significantly elevated in the cancer tissues.

Notable Publications

Author	Pubmed ID	Journal	Application
Bo Xu	35690146	J Biol Chem	WB
Fuji Yang	39921472	Adv Sci (Weinh)	WB,IF
S I Panpan	39496915	Mol Cell Biochem	WB,IHC


Storage

Store at -20°C. Stable for one year after shipment.


PBS with 0.02% sodium azide and 50% glycerol, pH7.3

Aliquoting is unnecessary for -20°C storage


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 66823-1-1g (OGT antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded human lung cancer tissue slide using 66823-1-Ig (OGT antibody) at dilution of 1:500 (under 10x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded human lung cancer tissue slide using 66823-1-Ig (OGT antibody) at dilution of 1:500 (under 40x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (4% PFA) fixed HepG2 cells using OGT antibody (66823-1-lg, Clone: 2B2A6) at dilution of 1:400 and CoraLite®488-Conjugated Goat Anti-Mouse IgG(H+L), CL594-phalloidin (red).