For Research Use Only

AGER Monoclonal antibody

Catalog Number:66833-1-lg 2 Publications



Purification Method:

CloneNo.:

IHC 1:150-1:600

3D4G11

Protein G purification

Recommended Dilutions: WB 1:1000-1:6000

Basic Information

Applications

Catalog Number: 66833-1-Ig

Size: 1500 µg/ml Source: Mouse

Isotype: IgG1 Immunogen Catalog Number:

AG9657

Tested Applications:

IHC, WB, ELISA
Cited Applications:

WB

Species Specificity: Human, Pig, Rat Cited Species: human, mouse

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

GenBank Accession Number:

BC020669 GeneID (NCBI):

UNIPROT ID: Q15109 Full Name:

> Calculated MW: 404 aa, 43 kDa Observed MW: 45 kDa

advanced glycosylation end productspecific receptor

Positive Controls:

WB: pig lung tissue, rat lung tissue

IHC: human lung tissue,

Background Information

Advanced glycosylation end product-specific receptor (AGER, also known as RAGE) is a multiligand receptor of the immunoglobulin superfamily. It interacts with distinct molecules implicated in homeostasis, development, and inflammation, and certain diseases such as diabetes and Alzheimer disease.

Notable Publications

Author	Pubmed ID	Journal	Application
Chengcheng Liu	34653793	Mol Immunol	WB
Lei Yang	34856357	Cell Signal	WB

Storage

Storage:


Store at -20°C. Stable for one year after shipment.

Storage Buffer


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


Selected Validation Data



pig lung tissue were subjected to SDS PAGE followed by western blot with 66833-1-lg (AGER antibody) at dilution of 1:3000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded human lung tissue slide using 66833-1-Ig (AGER antibody) at dilution of 1:300 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded human lung tissue slide using 66833-1-Ig (AGER antibody) at dilution of 1:300 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).