## For Research Use Only

## AGER Monoclonal antibody

Catalog Number:66833-1-lg 2 Publications



**Purification Method:** 

Protein G purification

Recommended Dilutions:

WB 1:1000-1:6000 IHC 1:150-1:600

CloneNo.:

3D4G11

Positive Controls:

IHC: human lung tissue,

WB: pig lung tissue, rat lung tissue

**Basic Information** 

Catalog Number: 66833-1-Ig

 $\begin{array}{lll} \text{Size:} & \text{GeneID (NCBI):} \\ \textbf{1500} \ \mu \, \textbf{g/ml} & \textbf{177} \\ \text{Source:} & \text{UNIPROT ID:} \end{array}$ 

Mouse Q15109

Isotype: Full Name:
IgG1 advanced glycosylation end product-

Immunogen Catalog Number:

AG9657

Applications

Tested Applications: IHC, WB, ELISA

Cited Applications:

WB

Species Specificity: Human, Pig, Rat Cited Species: human, mouse

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (\*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

**Background Information** 

Advanced glycosylation end product-specific receptor (AGER, also known as RAGE) is a multiligand receptor of the immunoglobulin superfamily. It interacts with distinct molecules implicated in homeostasis, development, and inflammation, and certain diseases such as diabetes and Alzheimer disease.

GenBank Accession Number:

BC020669

specific receptor

Calculated MW: 404 aa, 43 kDa Observed MW: 45 kDa

**Notable Publications** 

Author	Pubmed ID	Journal	Application
Chengcheng Liu	34653793	Mol Immunol	WB
Lei Yang	34856357	Cell Signal	WB

Storage

Storage:


Store at -20°C. Stable for one year after shipment.

Storage Buffer:


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


## **Selected Validation Data**



pig lung tissue were subjected to SDS PAGE followed by western blot with 66833-1-1g (AGER antibody) at dilution of 1:3000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded human lung tissue slide using 66833-1-Ig (AGER antibody) at dilution of 1:300 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded human lung tissue slide using 66833-1-Ig (AGER antibody) at dilution of 1:300 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).