For Research Use Only

AMPK Gamma 1 Monoclonal antibody

Catalog Number:67182-1-lg Featured Product



Basic Information

Catalog Number: 67182-1-lg

Size: 1000 µg/ml Source: Mouse Isotype: lgG1

Immunogen Catalog Number:

AG28907

WB, ELISA

Tested Applications:

Species Specificity: Human, mouse, rat

GenBank Accession Number:

BC000358 GeneID (NCBI): **UNIPROT ID:** P54619

protein kinase, AMP-activated, gamma 1 non-catalytic subunit

Calculated MW: 38 kDa Observed MW:

Full Name:

36-38 kDa

Positive Controls:

WB: HeLa cells, HEK-293 cells, HepG2 cells, Jurkat cells, K-562 cells, HSC-T6 cells, PC-12 cells, NIH/3T3

Purification Method:

CloneNo.:

1E11D4

Protein G purification

Recommended Dilutions:

WB 1:5000-1:20000

cells. RAW 264.7 cells

Background Information

Applications

Protein kinase, AMP-activated, gamma 1 non-catalytic subunit (PRKAG1, synonyms: AMPKG, MGC8666) is a regulatory subunit of the AMP-activated protein kinase (AMPK). AMPK is a heterotrimer consisting of an catalytic subunit, and non-catalytic and subunits. AMPK is an important energy-sensing enzyme that monitors cellular energy status. In response to cellular metabolic stresses, AMPK is activated, and thus phosphorylates and iAMP-activated protein kinase (AMPK) is a highly conserved heterotrimeric serine/threonine kinase widely characterised as a sensor of cellular energetic stress. AMPK is a heterotrimeric complex consisting of a catalytic α -subunit and two regulatory subunits (β and γ). AMPK is an important energy-sensing enzyme that monitors cellular energy status. In response to cellular metabolic stresses, AMPK is activated, and thus phosphorylates and inactivates acetyl-CoA carboxylase (ACC) and beta-hydroxy beta-methylglutaryl-CoA reductase (HMGCR), key enzymes involved in regulating de novo biosynthesis of fatty acid and cholesterol. AMPK gamma 1 is one of the gamma regulatory subunits of AMPK.

Storage


Storage:


Store at -20°C. Stable for one year after shipment.

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

Selected Validation Data





Various lysates were subjected to SDS PAGE followed by western blot with 67182-1-1g (AMPK gamma 1 antibody) at dilution of 1:20000 incubated at room temperature for 1.5 hours. The membrane was stripped and reblotted with HRP-conjugated Alpha Tubulin Monoclonal antibody (HRP-66031) as loading control.

WB result of AMPK Gamma 1 antibody (67182-1-lg; 1:20000; incubated at room temperature for 1.5 hours) with sh-Control and sh-AMPK Gamma 1 transfected HEK-293 cells.