

# ACO2 Monoclonal antibody

Catalog Number: 67509-1-Ig

## Basic Information

Catalog Number:	GenBank Accession Number:	Purification Method:
67509-1-Ig	BC014092	Protein G purification
Size:	GeneID (NCBI):	CloneNo.:
1600 µg/ml	50	1F1G4
Source:	UNIPROT ID:	Recommended Dilutions:
Mouse	Q99798	WB 1:5000-1:50000
Isotype:	Full Name:	IHC 1:500-1:2000
IgG1	aconitase 2, mitochondrial	
Immunogen Catalog Number:	Calculated MW:	
AG17784	85 kDa	
	Observed MW:	
	85 kDa	

## Applications

Tested Applications:	Positive Controls:
FC, IHC, WB, ELISA	WB : PC-12 cells, pig brain tissue, rat brain tissue, mouse brain tissue, HEK-293 cells, pig cerebellum tissue, rat cerebellum tissue, mouse cerebellum tissue, Neuro-2a cells, rabbit brain tissue
Species Specificity:	IHC : human liver cancer tissue,
Human, Mouse, Rat, Pig	
<b>Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0</b>	

## Background Information

ACO2(aconitate hydratase, mitochondrial) is also named as citrate hydro-lyase and belongs to the aconitase/IPM isomerase family. It plays a key function in cellular energy production, and loss of its activity has a major impact on cellular and organismal survival. Western blot shows two bands of 83 kDa and 40 kDa. The 40 kDa fragment decreases with age and oxidative stress, whereas other fragmentation products with molecular weights between 40 and 83 kDa increased with age and MnSOD(mitochondrial manganese superoxide dismutase) deficiency(PMID:12459471). Defects in ACO2 are the cause of infantile cerebellar-retinal degeneration (ICRD).

## Storage

**Storage:**  
Store at -20°C. Stable for one year after shipment.  
**Storage Buffer:**  
PBS with 0.02% sodium azide and 50% glycerol pH 7.3.  
Aliquoting is unnecessary for -20°C storage


For technical support and original validation data for this product please contact:

T: 4006900926


E: [Proteintech-CN@ptglab.com](mailto:Proteintech-CN@ptglab.com)W: [ptgcn.com](http://ptgcn.com)

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 67509-1-Ig (ACO2 antibody) at dilution of 1:20000 incubated at room temperature for 1.5 hours. The membrane was stripped and reblotted with HRP-conjugated Alpha Tubulin Monoclonal antibody (HRP-66031) as loading control.




Immunohistochemical analysis of paraffin-embedded human liver cancer tissue slide using 67509-1-Ig (ACO2 antibody) at dilution of 1:2000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).




Immunohistochemical analysis of paraffin-embedded human liver cancer tissue slide using 67509-1-Ig (ACO2 antibody) at dilution of 1:2000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



1X10<sup>6</sup> HeLa cells were intracellularly stained with 0.4 ug Anti-Human ACO2 (67509-1-Ig, Clone:1F1G4) and CoraLite®488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L) at dilution 1:1000 (red), or 0.4 ug Mouse IgG1 Isotype Control (MOPC-21) (65124-1-Ig, Clone: MOPC-21) (blue). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).



Various lysates were subjected to SDS PAGE followed by western blot with 67509-1-Ig (ACO2 antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.



HEK-293 cells were subjected to SDS PAGE followed by western blot with 67509-1-Ig (ACO2 antibody) at dilution of 1:20000 incubated at room temperature for 1.5 hours.