For Research Use Only

## ATP1B3 Monoclonal antibody

Catalog Number:67554-1-lg Featured Product

1 Publications




|                        | Catalog Number:<br>67554-1-lg   | GenBank Accession Number:<br>BC011835              | Purification Method:<br>Protein A purification  |  |
|------------------------|---|--|---|--|
|                        | Size:   | GenelD (NCBI):                                     | CloneNo.:   |  |
|                        | 1300 µg/ml  | 483  | 1H9C1   |  |
|                        | Source:   | UNIPROT ID:  | Recommended Dilutions:  |  |
|                        | Mouse   | P54709   | WB 1:1000-1:4000  |  |
|                        | Isotype:<br>IgG1<br>Immunogen Catalog Number:<br>AG30127  | Full Name:   | IHC 1:50-1:500  |  |
|                        |   | ATPase, Na+/K+ transporting, beta 3<br>polypeptide |   |  |
|                        |   | Calculated MW:                                     |   |  |
|                        |   | 32 kDa   |   |  |
|                        |   | Observed MW:<br>38-43 kDa                          |   |  |
| Applications           | Tested Applications:  | Positive   | Positive Controls:<br>WB : Jurkat cells, HeLa cells, HepG2 cells, human<br>placenta tissue, THP-1 cells |  |
|                        | IHC, WB, ELISA<br>Cited Applications:   |  |   |  |
|                        | IHC   |  |   |  |
|                        | Species Specificity:<br>Human   | IHC : human liver cancer tissue,                   |   |  |
|                        | Cited Species:<br>human   |  |   |  |
|                        | Note-IHC: suggested antige<br>TE buffer pH 9.0; (*) Alterno<br>retrieval may be performed<br>buffer pH 6.0  | atively, antigen                                   |   |  |
| Background Information | ATP1B3 (also known as CD298) is the $\beta$ 3 subunit of Na+/K+-ATPase which functions to maintain sodium and<br>potassium gradients across membranes involved in cellular activities. ATP1B3 is a glycosylated protein and there<br>are fully and intermediately glycosylated forms of ATP1B3 in mammalian cells. The predicted MW of ATP1B3 is<br>around 32 kDa, while various forms (38-43 kDa) can be observed due to the different level of glycosylation (PMID:<br>30792309, 16339171, 17176442). |  |   |  |
|                        | - ··· · · · · · · · · · · · · · · · · ·   |  |   |  |
| Notable Publications   | Author  | Pubmed ID Journal                                  | Application   |  |

For technical support and original validation data for this product please contact: E: Proteintech-CN@ptglab.com T: 4006900926 W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

## Selected Validation Data





Jurkat cells were subjected to SDS PAGE followed by western blot with 67554-1-1g (ATP1B3 antibody) at dilution of 1:2000 incubated at room temperature for 1.5 hours. Immunohistochemical analysis of paraffinembedded human liver cancer tissue slide using 67554-1-1g (ATP1B3 antibody) at dilution of 1:100 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).