For Research Use Only

ACADM Monoclonal antibody

Catalog Number:67742-1-lg 1 Publications




Basic Information	Catalog Number: 67742-1-lg	GenBank Accession Number: BC005377	Purification Method: Protein G purification	
	Size:	GeneID (NCBI):	CloneNo.:	
	1000 µg/ml	34	1A11G6	
	Source: Mouse	UNIPROT ID:	Recommended Dilutions:	
		P11310	WB 1:2000-1:10000 IHC 1:250-1:1000	
	Isotype: IgG1 Immunogen Catalog Number: AG30620	Full Name: THC 1.250-1.1000 acyl-Coenzyme A dehydrogenase, C-4 ^{IF} 1:200-1:800 to C-12 straight chain		
		Calculated MW:		
		421 aa, 47 kDa		
		Observed MW: 42 kDa		
Applications	Tested Applications:	Positive	Controls:	
	IF-P, IHC, WB, ELISA	WB:PC-3	cells, HSC-T6 cells, rat liver tissue, HeLa	
	Cited Applications: WB	cells, HEK-293 cells, HepG2 cells		
		IHC : hum	IHC : human liver tissue, human heart tissue	
	Species Specificity: Human, rat, pig	IF : huma	IF : human liver cancer tissue,	
	Cited Species: rat			
	Note-IHC: suggested antige. TE buffer pH 9.0; (*) Alterno retrieval may be performed buffer pH 6.0	atively, antigen		
Background Information	lengths of 4 to 16. It catalyzes the in ACADM are the cause of medium	reaction: Acyl-CoA + acceptor = 2,3-d	family. This enzyme is specific for acyl cha ehydroacyl-CoA + reduced acceptor. Defect ficiency (MCAD deficiency). This protein ca	
Background Information	lengths of 4 to 16. It catalyzes the in ACADM are the cause of mediu exsit as a dimer(PMID:8962055). T	reaction: Acyl-CoA + acceptor = 2,3-d m-chain acyl-CoA dehydrogenase det	ehydroacyl-CoA + reduced acceptor. Defect	


For technical support and original validation data for this product please contact: E: Proteintech-CN@ptglab.com T: 4006900926 W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 67742-1-Ig (ACADM antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours. The membrane was stripped and reblotted with HRP-conjugated GAPDH Monoclonal antibody (HRP-60004) as loading control.




rat liver tissue were subjected to SDS PAGE followed by western blot with 67742-1-1g (ACADM antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours.




Immunohistochemical analysis of paraffinembedded human liver tissue slide using 67742-1-Ig (ACADM antibody) at dilution of 1:500 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded human liver tissue slide using 67742-1-Ig (ACADM antibody) at dilution of 1:500 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (4% PFA) fixed human liver cancer tissue using ACADM antibody (67742-1-Ig, Clone: 1A11G6) at dilution of 1:400 and CoraLite®488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L).



HSC-T6 cells were subjected to SDS PAGE followed by western blot with 67742-1-Ig (ACADM antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours.