For Research Use Only

GSTK1 Recombinant antibody

Catalog Number:81527-1-RR



Basic Information

Catalog Number: GenBank Accession Number: 81527-1-RR BC050715

Protein A purification GeneID (NCBI): Size: CloneNo.: 250 µg/ml 373156 **UNIPROT ID:** Source:

Rabbit Q9Y2Q3 Full Name: Isotype: glutathione S-transferase kappa 1

Calculated MW: Immunogen Catalog Number:

AG6036 32 kDa

Observed MW: 27 kDa

Recommended Dilutions:

Purification Method:

WB 1:5000-1:50000 IHC 1:50-1:500 IF 1:200-1:800

Applications

Tested Applications: IF/ICC, IHC, WB, ELISA Species Specificity: human, mouse, rat

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate

buffer pH 6.0

Positive Controls:

WB: HEK-293 cells, HeLa cells, HepG2 cells, LO2 cells, Raji cells, Daudi cells

IHC: mouse testis tissue, mouse liver tissue, human pancreas cancer tissue, human lung cancer tissue, human liver cancer tissue, human stomach cancer tissue

IF: HeLa cells.

Background Information

 ${\sf GSTK1, glutathione S-transferase \, kappa \, 1, belongs \, to \, the \, superfamily \, of \, glutathione \, S-transferase \, ({\sf GST}), \, which \, is \, is \, in the superfamily \, of \, glutathione \, S-transferase \, ({\sf GST}), \, which \, is \, in the superfamily \, of \, glutathione \, S-transferase \, ({\sf GST}), \, which \, is \, in the superfamily \, of \, glutathione \, S-transferase \, ({\sf GST}), \, which \, is \, in the superfamily \, of \, glutathione \, S-transferase \, ({\sf GST}), \, which \, is \, in the superfamily \, of \, glutathione \, S-transferase \, ({\sf GST}), \, which \, in the superfamily \, of \, glutathione \, S-transferase \, ({\sf GST}), \, which \, in the superfamily \, of \, glutathione \, S-transferase \, ({\sf GST}), \, which \, in the superfamily \, of \, glutathione \, S-transferase \, ({\sf GST}), \, which \, in the superfamily \, of \, glutathione \, S-transferase \, ({\sf GST}), \, which \, in the superfamily \, of \, glutathione \, S-transferase \, ({\sf GST}), \, which \, in the superfamily \, of \, glutathione \, S-transferase \, ({\sf GST}), \, which \, in the superfamily \, of \, glutathione \, S-transferase \, ({\sf GST}), \, which \, in the superfamily \, of \, glutathione \, S-transferase \, ({\sf GST}), \, which \, in the superfamily \, of \, glutathione \, S-transferase \, ({\sf GST}), \, which \, in the superfamily \, of \, glutathione \, S-transferase \, ({\sf GST}), \, which \, in the superfamily \, of \, glutathione \, S-transferase \, ({\sf GST}), \, which \, in the superfamily \, of \, glutathione \, S-transferase \, ({\sf GST}), \, which \, in the superfamily \, of \, glutathione \, S-transferase \, ({\sf GST}), \, which \, in the superfamily \, of \, glutathione \, S-transferase \, ({\sf GST}), \, which \, in the superfamily \, of \, glutathione \, S-transferase \, ({\sf GST}), \, which \, in the superfamily \, of \, glutathione \, S-transferase \, ({\sf GST}), \, which \, in the superfamily \, of \, glutathione \, S-transferase \, ({\sf GST}), \, which \, in the superfamily \, of \, glutathione \, S-transferase \, ({\sf GST}), \, which \, in the superfamily \, of \, glutathione \, S-transferase \, ({\sf GST}), \, which \, in the superfamily \, of \,$ related to oxidative stress (PMID:16081649). GSTK1 expression is negatively correlated with obesity and hypertrophic cardiomyopathy (PMID:21428694, PMID:27378925). GSTK1 was expressed in the peroxisomal and mitochondrial fractions and localized to peroxisomes (PubMed:14742434).

Storage


Storage:

Store at -20°C. Stable for one year after shipment. Storage Buffer:


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 81527-1-RR (GSTK1 antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded mouse testis tissue slide using 81527-1-RR (GSTK1 antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (4% PFA) fixed Hela cells using GSTK1 antibody (81527-1-RR, Clone: 3J1) at dilution of 1:400 and Coralite®594-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).