## For Research Use Only

## Calretinin Recombinant antibody

Catalog Number:82811-1-RR



Basic Information	Catalog Number: 82811-1-RR	0		Purification Method: Protein A purification
	Concentration: 250 ug/ml	GenelD (NCBI): 794		CloneNo.: 2F20
	Source: Rabbit	UNIPROT ID: P22676 Full Name: calbindin 2 Calculated MW: 29 kDa		Recommended Dilutions: WB: 1:1000-1:5100 IHC: 1:50-1:500 IF/ICC: 1:125-1:500 FC (Intra): 0.25 ug per 10^6 cells in a 100 μl suspension
	Isotype: IgG			
	Immunogen Catalog Number: AG2924			
		Observed MW: 29 kDa		
Applications	Tested Applications: WB, IHC, IF/ICC, FC (Intra), ELISA Species Specificity: human, mouse, rat, pig		Positive Controls:	
			WB : SH-SY5Y cells, U2OS cells, fetal human brain tissue, mouse brain tissue, rat brain tissue, pig brain tissue	
	IF/ICC : SH-SY5Y cells,			
	FC (Intra) : SH-SY5Y cells,			
	Background Information	Calbindin 2 (calretinin), is an intracellular calcium-binding protein belonging to the troponin C superfamily. Members of this protein family have six EF-hand domains which bind calcium. This protein plays a role in diverse cellular functions, including message targeting and intracellular calcium buffering. It also functions as a modulator of neuronal excitability, and is a diagnostic marker for some human diseases, including Hirschsprung disease and some cancers. Calretinin is a useful marker for differentiating malignant mesothelioma from carcinomas.		
Storage	Storage: Store at -20°C. Stable for one year Storage Buffer: PBS with 0.02% sodium azide and Aliquoting is unnecessary for -20°	50% glycerol, pH7.3		

For technical support and original validation data for this product please contact:T: 4006900926E: Proteintech-CN@ptglab.comW: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

## Selected Validation Data



Immunofluorescent analysis of (4% PFA) fixed adult mouse cerebellum tissue using Calretinin antibody (82811-1-RR) at dilution of 1:500 and Alexa Fluor 594 AffiniPure Donkey Anti-Rabbit IgG (H+L). (Image produced by Watt Lab at McGill University)



Various lysates were subjected to SDS PAGE followed by western blot with 82811-1-RR (Calretinin antibody) at dilution of 1:2500 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded mouse brain tissue slide using 82811-1-RR (Calretinin antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded mouse brain tissue slide using 82811-1-RR (Calretinin antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (4% PFA) fixed SH-SY5Y cells using Calretinin antibody (82811-1-RR, Clone: 2F20) at dilution of 1:250 and Multi-rAb CoraLite ® Plus 594-Goat Anti-Rabbit Recombinant Secondary Antibody (H+L) (RGAR004).



Immunohistochemical analysis of paraffinembedded mouse brain tissue slide using 82811-1-RR (Calretinin antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



1x10^6 SH-SY5Y cells were intracellularly stained with 0.25 ug Calretinin Recombinant antibody (82811-1-RR, Clone:2F20) and CoraLite@488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L) (SA00013-2)(red), or 0.25 ug Isotype Control (blue). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).