For Research Use Only

IFT88 Recombinant antibody

Catalog Number:82886-1-RR



Basic Information

Catalog Number: 82886-1-RR

Size: 1000 μ g/ml

Source: Rabbit

Isotype:

BC030776

GeneID (NCBI):
8100

UNIPROT ID:

Q13099

Full Name:
intraflagellar transport 88 homolog

GenBank Accession Number:

Immunogen Catalog Number: (Chlamydomonas)
AG4980 Calculated MW:

94 kDa Observed MW: 88-95 kDa

Applications

Tested Applications: IHC, WB, ELISA Species Specificity: Human, mouse, rat

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0 Purification Method: Protein A purification

230074F8

CloneNo.:

Recommended Dilutions: WB 1:2000-1:10000 IHC 1:250-1:1000

Positive Controls:

WB: HEK-293 cells, NIH/3T3 cells, human testis tissue, mouse testis tissue, rat testis tissue

IHC: mouse heart tissue,

Background Information

Intraflagellar transport (IFT), mediated by molecular motors and IFT particles, is an important transport process that occurs in the cilium and has been shown to be essential for the assembly and maintenance of cilia and flagella in many organisms. IFT88 (intraflagellar transport protein 88; also known as TG737 or TTC10) is a component of IFT particles and required for cilium biogenesis. Defects in IFT88/Tg737 lead to polycystic kidney disease (11062270). IFT88 localizes to spindle poles during mitosis and is required for spindle orientation in mitosis (21441926). This antibody was raised against the C-terminal region of human IFT88 and can detect the endogenous level of IFT88.

Storage


Storage:

Store at -20°C. Stable for one year after shipment. Storage Buffer:


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

Selected Validation Data



Immunohistochemical analysis of paraffinembedded mouse heart tissue slide using 82886-1-RR (IFT88 antibody) at dilution of 1:500 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Various lysates were subjected to SDS PAGE followed by western blot with 82886-1-RR (IFT88 antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours.