For Research Use Only

Atlantic Blue™ Anti-Human CD107a / LAMP1 (H4A3)

Proteintech

Antibodies | ELISA kits | Proteins

www.ptglab.com

Catalog Number: AB-65051

Basic Information

Catalog Number:

AB-65051

Size:

100tests, 5 μ l/test

Source: Mouse

Isotype: IgG1, kappa

P11279 Full Name:

lysosomal-associated membrane

GenBank Accession Number:

protein 1

BC006345

3916

GeneID (NCBI):

ENSEMBL Gene ID:

ENSG00000185896

UNIPROT ID:

Purification Method:

Protein G purification

CloneNo.: H4A3

Excitation/Emission maxima

wavelengths: 404 nm / 458 nm

Applications

Tested Applications:

Species Specificity:

Human

FC (Intra)

Background Information

LAMP1 (CD107a) is a heavily glycosylated membrane protein enriched in the lysosomal membrane. LAMP1 is extensively glycosylated with asparagine-linked oligosaccharides which protect it from intracellular proteolysis (PMID: 10521503). Although LAMP1 is expressed largely in the endosome-lysosomal membrane of cells, it is also found on the plasma membrane (PMID: 16168398). Elevated LAMP1 expression at the cell surface has also been detected during platelet and granulocytic cell activation, as well as in some tumor cells (PMID: 29085473). LAMP1 functions to provide selectins with carbohydrate ligands. This protein has also been shown to be a marker of degranulation on lymphocytes such as CD8+ and NK cells and may also play a role in tumor cell differentiation and metastasis (PMID: 18835598; 29085473; 9426697).

Storage


Storage:

Store at 2-8 °C. Avoid exposure to light. Stable for one year after shipment.


Storage Buffer:

PBS with 0.09% sodium azide and 0.5% BSA.

Selected Validation Data



1X10^6 HeLa cells were intracellularly stained with 5 ul Atlantic Blue™ Anti-Human LAMP1 (AB-65051, Clone:H4A3). Cells were fixed with 4% PFA and permeabilized with Intracellular Staining Permeabilization Wash Buffer.



1X10^6 HeLa cells were intracellularly stained with 5 ul Atlantic Blue™ Anti-Human LAMP1 (AB-65051, Clone:H4A3). Cells were fixed with 4% PFA and permeabilized with Intracellular Staining Permeabilization Wash Buffer.