## For Research Use Only

## APC Anti-Mouse CD8a (53-6.7)


Catalog Number: APC-65069 5 Publications



<b>Basic Information</b>	Catalog Number: APC-65069	GenBank Ad BC030679	ccession Number:	Purification Method: Affinity purification	
	Size: 100ug, 0.1 mg/ml	Genel D (NC 12525	BI):	CloneNo.: 53-6.7	
	Source: Rat	UNIPROT IE P01731	):	Excitation/Emission maxima wavelengths:	
	Isotype: Full Name: IgG2a, kappa CD8 antigen, alpha chain			650 nm / 660 nm	
Applications	Tested Applications: FC				
	Cited Applications: IF, FC				
	Species Specificity: Mouse				
	Cited Species: mouse				
Background Information	CD8 a or as a heterodimer majority of class I-restricte have been found on some g (PMID: 2111591; 1831127; 1	of CD8 a and CD8 β (P ed T cells express mostl gut intraepithelial T cel 8420975). CD8 acts as a	MID: 3264320; 825379 y the CD8 $\alpha \beta$ heteroo lls , on some T cell rec a co-receptor that bind	d chains. It can be present as a homodimer of p1). CD8 is found on most thymocytes. The dimer while CD8 $\alpha$ $\alpha$ homodimers alone eptor (TCR) $\gamma$ $\delta$ T cells and on NK cells s to MHC class-l and participates in cytotoxi quired for positive selection of CD4-/CD8+T	
	CD8 a or as a heterodimer majority of class I-restricte have been found on some g (PMID: 2111591; 1831127; T cell activation (PMID: 849	of CD8 a and CD8 β (P ed T cells express mostl gut intraepithelial T cel 8420975). CD8 acts as a	MID: 3264320; 825379 y the CD8 $\alpha \beta$ heteroo lls , on some T cell rec a co-receptor that bind	D1). CD8 is found on most thymocytes. The dimer while CD8 $\alpha \alpha$ homodimers alone eptor (TCR) $\gamma \delta$ T cells and on NK cells s to MHC class-I and participates in cytotoxi	
	CD8 a or as a heterodimer majority of class I-restricte have been found on some g (PMID: 2111591; 1831127; T cell activation (PMID: 849 cells (PMID: 1968084).	of CD8 a and CD8 β (P d T cells express mostl gut intraepithelial T cel 8420975). CD8 acts as a 99079). During T cell de	MID: 3264320; 825379 y the CD8 α β heteroo ils , on some T cell reco a co-receptor that bind evelopment, CD8 is rec	D1). CD8 is found on most thymocytes. The dimer while CD8 $\alpha \alpha$ homodimers alone eptor (TCR) $\gamma \delta$ T cells and on NK cells s to MHC class-I and participates in cytotoxic quired for positive selection of CD4-/CD8+T	
	CD8 a or as a heterodimer majority of class I-restricte have been found on some g (PMID: 2111591; 1831127; T cell activation (PMID: 849 cells (PMID: 1968084).	of CD8 α and CD8 β (P d T cells express mostl gut intraepithelial T cel 8420975). CD8 acts as a 99079). During T cell de Pubmed ID	MID: 3264320; 825379 y the CD8 α β heteroo lls, on some T cell reca a co-receptor that bind evelopment, CD8 is rec Journal	D1). CD8 is found on most thymocytes. The dimer while CD8 a a homodimers alone eptor (TCR) γ δ T cells and on NK cells s to MHC class-I and participates in cytotoxia quired for positive selection of CD4-/CD8+T Application	
Background Information	CD8 a or as a heterodimer majority of class I-restricte have been found on some g (PMID: 2111591; 1831127; 1 T cell activation (PMID: 849 cells (PMID: 1968084). Author Dingyi Wang	of CD8 a and CD8 β (P d T cells express mostl gut intraepithelial T cel 8420975). CD8 acts as a 99079). During T cell de <b>Pubmed ID</b> 35236203	MID: 3264320; 825379 y the CD8 α β heteroo ils , on some T cell reco a co-receptor that bind evelopment, CD8 is rec Journal Drug Deliv	p1). CD8 is found on most thymocytes. The dimer while CD8 α α homodimers alone eptor (TCR) γ δ T cells and on NK cells s to MHC class-I and participates in cytotoxi quired for positive selection of CD4-/CD8+ T Application FC	

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

## Selected Validation Data



1X10^6 mouse splenocytes were surface co-stained with CoraLite® Plus 488 Anti-Mouse CD3 and 0.2 ug APC Anti-Mouse CD8a (APC-65069, Clone:53-6.7) or 0.2 ug Isotype Control. Cells were not fixed.