For Research Use Only

CoraLite® Plus 488-conjugated BST2 Polyclonal antibody



Purification Method:

IF 1:200-1:800

wavelengths: 493 nm / 522 nm

Antigen affinity purification

Excitation/Emission maxima

Recommended Dilutions:

Catalog Number: CL488-13560

Featured Product

Basic Information

Applications

Catalog Number: CL488-13560

1000 μg/ml

Source: Rabbit Isotype:

Immunogen Catalog Number:

mmunogen Catalog Number: AG4430

G4430

Tested Applications: IF/ICC
Species Specificity:

human, mouse

GenBank Accession Number:

BC033873 GeneID (NCBI):

UNIPROT ID:

Q10589 Full Name:

Calculated MW:

bone marrow stromal cell antigen 2

180 aa, 20 kDa Observed MW: 30-36 kDa, 50-70 kDa

Positive Controls: IF: HepG2 cells,

Background Information

BST2, also named as CD317 and Tetherin, belongs to the tetherin family. It may be involved in the sorting of secreted proteins and it is involved in pre-B-cell growth. BST2 is an antiretroviral defense protein, that blocks release of retrovirus from the cell surface. Depleted unpon HIV-1 infection by viral VPU protein through 20S proteasome degradation. Depleted upon infection by human Kaposi's sarcoma-associated herpesvirus (KSHV) through ubiquitination and subsequent degradation. BST2 may play a role in B-cell activation in rheumatoid arthritis. It is recently identified interferon-induced cellular proteins that restrict infections by retroviruses and filoviruses and of influenza virus and flaviviruses, respectively. BST2 is a plasma membrane proteins, tetherin inhibits virion particle release from infected cells. BST2 is effective against retroviruses and flavoviruses whilst IFITMs disrupt influenza and flavivirus infection. Observed MW of BST2 is 30-36 kDa (PMID: 19196977; 21237475).

Storage


Storage:

Store at -20°C. Avoid exposure to light. Stable for one year after shipment. Storage Buffer:

PBS with 50% Glycerol, 0.05% Proclin300, 0.5% BSA, pH 7.3.

Aliquoting is unnecessary for -20°C storage

Selected Validation Data



Immunofluorescent analysis of (4% PFA) fixed HepG2 cells using CoraLite® Plus 488 BST2 antibody (CL488-13560) at dilution of 1:400.