For Research Use Only

CoraLite® Plus 488-conjugated ATG16L1 Polyclonal antibody



Purification Method:

IF 1:50-1:500

wavelengths: 493 nm / 522 nm

Antigen affinity purification

Excitation/Emission maxima

Recommended Dilutions:

Catalog Number: CL488-29445

Basic Information

Catalog Number: CL488-29445

1000 µg/ml Source: Rabbit Isotype:

Immunogen Catalog Number:

AG31318

Tested Applications:

Species Specificity: Human, mouse, rat

GenBank Accession Number:

BC000061 GeneID (NCBI): 55054 **UNIPROT ID:**

Q676U5

607 aa, 68 kDa Observed MW: 68-75 kDa

Full Name: ATG16 autophagy related 16-like 1 (S.

cerevisiae) Calculated MW:

Positive Controls:

IF: HEK-293 cells,

Background Information

Human ATG16L1 is a 607 amino acid protein (~68 kDa) comprising three major domains: the N - terminal ATG5 binding domain (ATG5 - BD), the central coiled - coil domain (CCD) and a predicted C - terminal WD40 - domain. ATG16L1 α and β (Atg16L1 α , 63 kDa; and Atg16L1 β , 71 kDa) are the major isoforms expressed in intestinal epithelium and macrophages, and all isoforms encode exon 9, which contains Thr 300. Atg16L1 mediates the cellular degradative process of autophagy and is considered a critical regulator of inflammation based on its genetic association with inflammatory bowel disease. ATG16L1 has been implicated in Crohn's disease. (PMID: 24553140, PMID: 22740627, PMID: 28685931)

Storage


Applications

Store at -20°C. Avoid exposure to light. Stable for one year after shipment.

PBS with 50% Glycerol, 0.05% Proclin300, 0.5% BSA, pH 7.3.

Aliquoting is unnecessary for -20°C storage

Selected Validation Data



Immunofluorescent analysis of (-20°C Ethanol) fixed HEK-293 cells using Coralite® Plus 488 ATG16L1 antibody (CL488-29445) at dilution of 1:200.