For Research Use Only

## CoraLite® Plus 488 Anti-Mouse CD11c (N418)



Catalog Number: CL488-65130

**Basic Information** 

Catalog Number: CL488-65130

BC167225

Purification Method: Affinity purification

Ci---

GeneID (NCBI):

GenBank Accession Number:

CloneNo.:

100ug, 0.5 mg/ml

Armenian Hamster

16411

N418

Source:

UNIPROT ID: Q9QXH4 Recommended Dilutions: IF 1:200-1:800

Isotype:

Full Name:

Excitation/Emission maxima

integrin alpha X

wavelengths: 493 nm / 522 nm

**Applications** 

**Tested Applications:** 

FC, IF/ICC

Positive Controls:

IF: mouse splenocytes,

Species Specificity:

Mouse

**Background Information** 

Integrins are cell adhesion receptors that are heterodimers composed of non-covalently associated  $\alpha$  and  $\beta$  subunits (PMID: 9779984). CD11c, also known as integrin  $\alpha$  X, is a type I transmembrane glycoprotein present on a variety of cells, including monocytes/macrophages, granulocytes, a subset of B cells, NK cells and dendritic cells (PMID: 2897326; 1680915; 1694698; 17389580). As a result of its high level of expression on most dendritic cells, CD11c is typically considered to be a marker of conventional dendritic cells (PMID: 27119555). CD11c forms an  $\alpha$  /  $\beta$  heterodimer with CD18 (integrin  $\beta$  2). CD11c/CD18 acts a receptor for fibrinogen and is important in monocyte adhesion and chemotaxis (PMID: 1671533).

Storage


Storage:

Store at 2-8°C. Avoid exposure to light. Stable for one year after shipment.


Storage Buffer:

PBS with 0.1% sodium azide and 0.5% BSA, pH 7.3.

## Selected Validation Data



1X10^6 mouse splenocytes were surface co-stained with APC Anti-Mouse CD45R (B220) (RA3-6B2) and 0.25 ug CoraLite® Plus 488 Anti-Mouse CD11c (CL488-65130, Clone: N418) or 0.25 ug isotype control. Cells were not fixed.



Immunofluorescent analysis of mouse splenocytes using CoraLite® Plus 488 CD11c antibody (CL488-65130, Clone: N418) at dilution of 1:400.