CoraLite® Plus 488-conjugated 6*His, His-Tag Monoclonal antibody




Catalog Number:CL488-66005 <u>3 Publications</u>

Basic Information	Catalog Number: CL488-66005	GenBank Acc GeneID (NCB	ession Number: 81):	Purification Method: Protein G purification	
	Size:	Full Name:		CloneNo.:	
	1000 µg/ml Source:	Calculated M	1W:	1B7G5 Recommended Dilutions:	
	Mouse	1 kDa		IF 1:50-1:500	
	Isotype: IgG1			Excitation/Emission maxima wavelengths: 493 nm / 522 nm	
Applications	Tested Applications:		Positive Controls:		
	IF/ICC Cited Applications: FC	Cited Applications:			
	Cited Species: human				
	Protein tags are protein or peptide sequences located either on the C- or N- terminal of the target protein, which facilitates one or several of the following characteristics: solubility, detection, purification, localization and expression. His-tag is often used for affinity purification and binding assays. Expressed His-tagged proteins can be purified and detected easily because the string of histidine residues binds to several types of immobilized metal ions, including nickel, cobalt and copper, under specific buffer conditions. The His-tag antibody is a useful tool for monitoring of the His-tagged proteins, and recognizes His-tags placed at N-terminal, C-terminal, and internal regions of fusion proteins expressed in bacteria, insect, and mammalian cells.				
Background Informatio	expression. His-tag is often purified and detected easil ions, including nickel, coba monitoring of the His-tagge	f the following character nused for affinity purifica y because the string of h It and copper, under spe ed proteins, and recogniz	ristics: solubility, det ation and binding as: nistidine residues bin cific buffer condition zes His-tags placed a	tection, purification, localization and says. Expressed His-tagged proteins can b nds to several types of immobilized metal ns. The His-tag antibody is a useful tool for nt N-terminal, C-terminal, and internal	
	expression. His-tag is often purified and detected easil ions, including nickel, coba monitoring of the His-tagge	f the following character nused for affinity purifica y because the string of h It and copper, under spe ed proteins, and recogniz	ristics: solubility, det ation and binding as: nistidine residues bin cific buffer condition zes His-tags placed a	tection, purification, localization and says. Expressed His-tagged proteins can be nds to several types of immobilized metal ns. The His-tag antibody is a useful tool for nt N-terminal, C-terminal, and internal	
	expression. His-tag is often purified and detected easil ions, including nickel, coba monitoring of the His-tagge regions of fusion proteins e	f the following character n used for affinity purifice y because the string of h It and copper, under spe ed proteins, and recogniz expressed in bacteria, ins	ristics: solubility, det ation and binding as: istidine residues bin cific buffer condition zes His-tags placed a sect, and mammaliar	tection, purification, localization and says. Expressed His-tagged proteins can be nds to several types of immobilized metal ns. The His-tag antibody is a useful tool for at N-terminal, C-terminal, and internal n cells. Application	
	expression. His-tag is often purified and detected easil ions, including nickel, coba monitoring of the His-tagge regions of fusion proteins e	f the following character used for affinity purific y because the string of h It and copper, under spe ed proteins, and recogniz expressed in bacteria, ins Pubmed ID	ristics: solubility, det ation and binding as: nistidine residues bin cific buffer condition zes His-tags placed a sect, and mammaliar Journal	tection, purification, localization and says. Expressed His-tagged proteins can be nds to several types of immobilized metal ns. The His-tag antibody is a useful tool for at N-terminal, C-terminal, and internal n cells. Application	
Background Informatio	Author Jingwei Jiang	f the following character a used for affinity purific y because the string of h It and copper, under spe ed proteins, and recogniz expressed in bacteria, ins Pubmed ID 34998985	ristics: solubility, det ation and binding ass istidine residues bin cific buffer condition zes His-tags placed a sect, and mammaliar Journal Fish Shellfish Imr	tection, purification, localization and says. Expressed His-tagged proteins can be nds to several types of immobilized metal ns. The His-tag antibody is a useful tool for it N-terminal, C-terminal, and internal n cells. Application munol FC	

For technical support and original validation data for this product please contact: T: 4006900926 E: Proteintech-CN@ptglab.com W: ptgcn.com This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



Immunofluorescent analysis of (-20°C Ethanol) fixed Transfected HEK-293 cells using the CoraLite® Plus 488-conjugated version of this antibody, CL488-66005 (6°His, His-Tag antibody), at dilution of 1:100.