For Research Use Only

CoraLite® Plus 488-conjugated VEGFA Monoclonal antibody

Catalog Number:CL488-66828

Featured Product




Basic Information	Catalog Number: CL488-66828	GenBank Accession Number: BC065522	Purification Method: Protein A purification
	Size: 1000 µg/ml	GeneID (NCBI): 7422	CloneNo.: 2E2H9
	Source: Mouse	UNIPROT ID:Excitation/Emission maxinP15692wavelengths:Full Name:493 nm / 522 nmvascular endothelial growth factor A	Excitation/Emission maxima wavelengths:
	Isotype: IgG2b Immunogen Catalog Number: AG13500		493 nm / 522 nm
		Calculated MW: 412 aa, 46 kDa	
		Observed MW: 43 kDa	
Applications	Tested Applications: FC (Intra)		
	Species Specificity: Human		
Background Information	VEGFA, also named as VEGF or VPF, belongs to the PDGF/VEGF growth factor family. It is a growth factor active in angiogenesis, vasculogenesis and endothelial cell growth. VEGFA induces endothelial cell proliferation, promotes cell migration, inhibits apoptosis and induces permeabilization of blood vessels. It binds to the FLT1/VEGFR1 and KDR/VEGFR2 receptors, heparan sulfate and heparin. Defects in VEGFA are associated with microvascular complications of diabetes type 1 (MVCD1). VEGFA has 17 isoforms with MW from 16 to 45 kDa. Some isoforms have homodimer forms (e.g.; VEGFA189 38kDa or VEFGA110 34kDa). VEGF-A exists in at least seven homodimeric isoforms. The monomers consist of 121, 145, 148, 165, 183, 189, or 206 amino acids (PMID:15602010). This antibody can recognize all VEGFA isoforms.		
Storage	Storage: Store at -20°C. Avoid exposure to light. Stable for one year after shipment. Storage Buffer: PBS with 50% Glycerol, 0.05% Proclin300, 0.5% BSA, pH 7.3. Aliquoting is unnecessary for -20°C storage		

For technical support and original validation data for this product please contact:T: 4006900926E: Proteintech-CN@ptglab.comW: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



1X10^6 MCF-7 cells were intracellularly stained with 0.4 ug CoraLite® Plus 488 Anti-Human VEGFA (CL488-66828, Clone:2E2H9) (red), or 0.4 ug Control Antibody. Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).