For Research Use Only

CoraLite® Plus 488-conjugated U2AF2/U2AF65 Monoclonal antibody



Purification Method:

Catalog Number: CL488-68166

Basic Information

Catalog Number: GenBank Accession Number: CL488-68166 BC008740

 CL488-68166
 BC008740
 Protein A purification

 Size:
 GeneID (NCBI):
 CloneNo.:

 1000 μ g/ml
 11338
 1B7E7

Source: UNIPROT ID: Recommended Dilutions:

Mouse P26368 IF 1:50-1:500

 Isotype:
 Full Name:
 Excitation/Emission maxima

 IgG2a
 U2 small nuclear RNA auxiliary factor wavelengths:

Immunogen Catalog Number: 2 493 nm / 522 nm

AG8070 Calculated MW:

54 kDa Observed MW: 60-65 kDa

Applications

Tested Applications: Positive Controls:
FC (Intra), IF/ICC IF: HEK-293 cells,

Species Specificity: Human, rat, mouse

Background Information

U2AF2, also named as U2 auxiliary factor 65 kDa subunit, is a 475 amino acid protein, which belongs to the splicing factor SR family. U2AF2 is Necessary for the splicing of pre-mRNA. By recruiting PRPF19 and the PRP19C/Prp19 complex/NTC/Nineteen complex to the RNA polymerase II C-terminal domain (CTD), pre-mRNA, may couple transcription to splicing. The calculated molecular weight of U2AF2 is 53 kDa, but modified U2AF2 is about 65 kDa.

Storage


Storage:

Store at -20°C. Avoid exposure to light. Stable for one year after shipment.


PBS with 50% Glycerol, 0.05% Proclin300, 0.5% BSA, pH 7.3.

Aliquoting is unnecessary for -20°C storage

Selected Validation Data



Immunofluorescent analysis of (4% PFA) fixed HEK-293 cells using Coralite® Plus 488 U2AF2 antibody (CL488-68166, Clone: 1B7E7) at dilution of 1:100, CL594-phalloidin (red).



1X10^6 HEK-293 cells were intracellularly stained with 0.8 ug CoraLite® Plus 488 Anti-Human U2AF2 (CL488-68166, Clone:1B7E7) (red), or 0.8 ug CoraLite® Plus 488 Mouse IgG2a Isotype Control (C1.18.4) (CL488-65208, Clone: C1.18.4) (blue). Cells were fixed and permeabilized with Transcription Factor Staining Buffer Kit (PF00011).