

For Research Use Only

CoraLite® Plus 555-conjugated NANOG Polyclonal antibody



Catalog Number: CL555-14295

Featured Product

Basic Information

Catalog Number:

CL555-14295

Size:

1000 µg/mL

Source:

Rabbit

Isotype:

IgG

Immunogen Catalog Number:

AG5645

GenBank Accession Number:

BC160187

GeneID (NCBI):

79923

UNIPROT ID:

Q9H9S0

Full Name:

Nanog homeobox

Calculated MW:

35 kDa

Observed MW:

35-40 kDa

Purification Method:

Antigen affinity purification

Excitation/Emission maxima
wavelengths:

554 nm / 570 nm

Applications

Tested Applications:

FC (Intra)

Species Specificity:

human, mouse, rat

Background Information

Nanog is a member of the homeobox family of DNA binding transcription factors and has been shown to maintain embryonic stem (ES) cell self-renewal independently of leukemia inhibitory factor (LIF)/Stat3. Nanog mRNA is present in pluripotent mouse and human cell lines, and absent from differentiated cells. Functionally, Nanog works together with other key pluripotent factors (Oct4, Sox2, and Lin28) to reprogram human fibroblasts and generate induced pluripotent stem (iPS) cells. These key factors form a regulatory network to support or limit each other's expression level, which maintains the properties of ES cells. Affinity purified rabbit anti-Nanog can be used to demonstrate pluripotency of ES and IPS cells. There are two kinds of variants could be recognized by NANOG, one is normal form (~39kd), the other is post-translation modified form (~48kd) (21136380). Nanog exists two isoforms with molecular weight 34.4 kDa and 31.9 kDa. (PMID: 21969378)

Storage

Storage:

Store at -20°C. Avoid exposure to light. Stable for one year after shipment.

Storage Buffer:

PBS with 50% Glycerol, 0.05% Proclin300, 0.5% BSA, pH 7.3.

Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com

W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



1X10⁶ NCCIT cells were intracellularly stained with 0.2 ug CoraLite® Plus 555 Anti-Human NANOG (CL555-14295) (red), or 0.2 ug CoraLite® Plus 555 Rabbit IgG Control Antibody (CL555-30000) (Blue). Cells were fixed and permeabilized with Transcription Factor Staining Buffer Kit (PF00011).