

For Research Use Only

CoraLite®594-conjugated Ataxin 2 Polyclonal antibody



Catalog Number: CL594-21776

Featured Product

Basic Information

Catalog Number:

CL594-21776

Size:

1000 µg/ml

Source:

Rabbit

Isotype:

IgG

Immunogen Catalog Number:

AG16470

GenBank Accession Number:

BC114546

GeneID (NCBI):

6311

UNIPROT ID:

Q99700

Full Name:

ataxin 2

Calculated MW:

1313 aa, 140 kDa

Observed MW:

140-150 kDa

Purification Method:

Antigen affinity purification

Recommended Dilutions:

IF 1:50-1:500

Excitation/Emission maxima
wavelengths:

588 nm / 604 nm

Applications

Tested Applications:

FC (Intra), IF-P

Species Specificity:

human, mouse, rat

Positive Controls:

IF : mouse brain tissue,

Background Information

ATXN2 contains a repeat structure with 22 or 23 triplets coding for glutamine and the (CAG)₈CAA(CAG)₄CAA(CAG)₈ sequence; expansion of this domain to a size ≥ 34 triplets with a pure CAG sequence primarily causes autosomal dominant SCA2 [PMID:18418684], while ATXN2 expansions with CAA interruptions were observed as the cause of Levo-dopa responsive Parkinson's disease [PMID:10668726]. ATXN2 expansions associated with ALS were reported to be interrupted by at least one CAA triplet [PMID:21537950].

Storage

Storage:

Store at -20°C. Avoid exposure to light. Stable for one year after shipment.

Storage Buffer:

PBS with 50% Glycerol, 0.05% Proclin300, 0.5% BSA, pH 7.3.

Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com


W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



Immunofluorescent analysis of (4% PFA) fixed mouse brain tissue using CoraLite® 594 Ataxin 2 antibody (CL594-21776) at dilution of 1:200.



1X10⁶ HeLa cells were intracellularly stained with 0.4 ug CoraLite® 594 Anti-Human Ataxin 2 (CL594-21776) (red), or 0.4 ug Control Antibody. Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).