For Research Use Only

CoraLite®594-conjugated Beta Actin Monoclonal antibody

Proteintech®
Antibodies | ELISA kits | Proteins
www.ptglab.com

Catalog Number: CL594-60008

Basic Information

Catalog Number: GenBank Accession Number: CL594-60008 BC002409

CL594-60008 BC002409
Size: GeneID (NCBI):

Size: GeneID (NCBI): 1000 μ g/ml 60

Source: UNIPROT ID:

Mouse P60709

Isotype: Full Name:
IgM actin, beta

Immunogen Catalog Number: Calculated MW:

G0297 375 aa, 42 kDa

Purification Method:

Caprylic acid/ammonium sulfate

precipitation

CloneNo.: 7D2C10

Recommended Dilutions:

WB 1:1000-1:4000 IF 1:200-1:1000

Excitation/Emission maxima

wavelengths: 588 nm / 604 nm

Applications

Tested Applications:

IF/ICC, WB

Species Specificity:

human, mouse, rat, plant, Zebrafish, canine

Positive Controls:

WB: HepG2 cells, HeLa cells, HEK-293 cells

IF: HeLa cells, MDCK cells

Background Information

Beta actin, also named as ACTB and F-Actin, belongs to the actin family. Actins are highly conserved globular proteins that are involved in various types of cell motility and are ubiquitously expressed in all eukaryotic cells. At least six isoforms of actins are known in mammals and other vertebrates: alpha (ACTC1, cardiac muscle 1), alpha 1 (ACTA1, skeletal muscle) and 2 (ACTA2, aortic smooth muscle), beta (ACTB), gamma 1 (ACTG1) and 2 (ACTG2, enteric smooth muscle). Beta and gamma 1 are two non-muscle actin proteins. Most actins consist of 376aa, while ACTG2 (rich in muscles) has 375aa and ACTG1(found in non-muscle cells) has only 374aa. Beta actin has been widely used as the internal control in RT-PCR and Western Blotting as a 42-kDa protein. However, the 37-40 kDa cleaved fragment of beta actin can be generated during apoptosis process. This antibody is conjugated with CL594,Ex/Em 593 nm/614 nm.

Storage

Storage:


Store at -20°C. Avoid exposure to light. Stable for one year after shipment.

Storage Buffer


PBS with 50% Glycerol, 0.05% Proclin300, 0.5% BSA, pH 7.3.

Aliquoting is unnecessary for -20°C storage

Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with CL594-60008 (Beta Actin antibody) at dilution of 1:2000 incubated at room temperature for 1.5 hours.



Immunofluorescent analysis of (-20°C Methanol) fixed HeLa cells using CoraLite®594 Beta Actin antibody (CL594-60008, Clone: 7D2C10) at dilution of 1:200.