For Research Use Only

## CoraLite® Plus 647-conjugated TBP Monoclonal antibody



**Purification Method:** 

Protein A purification

Recommended Dilutions:

Excitation/Emission maxima

CloneNo.:

IF 1:50-1:500

wavelengths: 654 nm / 674 nm

2H3B2

Catalog Number: CL647-66166

**Basic Information** 

Catalog Number:

 CL647-66166
 BC110341

 Size:
 GeneID (NCBI):

 1000 μg/ml
 6908

 Source:
 UNIPROT ID:

 Mouse
 P20226

 Isotype:
 Full Name:

IgG2a TATA box binding protein
Immunogen Catalog Number: Calculated MW:

AG12383 338 aa, 38 kDa
Observed MW:

mouse/rat 33-36 kDa and human 37-

GenBank Accession Number:

43kDa

**Applications** 

Tested Applications: FC (Intra), IF/ICC

Species Specificity: human, mouse, rat, pig **Positive Controls:** 

IF: A431 cells,

## **Background Information**

The TATA binding protein (TBP) is a transcription factor that binds specifically to a DNA sequence TATA box. This DNA sequence is found about 25-30 base pairs upstream of the transcription start site in some eukaryotic gene promoters. TBP, along with a variety of TBP-associated factors, make up the TFIID, a general transcription factor that in turn makes up part of the RNA polymerase II preinitiation complex. As one of the few proteins in the preinitation complex that binds DNA in a sequence-specific manner, it helps position RNA polymerase II over the transcription start site of the gene. However, it is estimated that only 10-20% of human promoters have TATA boxes. Therefore, TBP is probably not the only protein involved in positioning RNA polymerase II. This antibody detects human TBP (-40 kDa) and mouse/rat Tbp (-35 kDa).

## Storage


Storage:


Store at -20°C. Avoid exposure to light. Stable for one year after shipment. Storage Buffer:

PBS with 50% Glycerol, 0.05% Proclin300, 0.5% BSA, pH 7.3.

Aliquoting is unnecessary for -20°C storage

## Selected Validation Data







Immunofluorescent analysis of (4% PFA) fixed A431 cells using CoraLite® Plus 647 TBP antibody (CL647-66166, Clone: 2H3B2) at dilution of 1:200.