

# CoraLite® Plus 647-conjugated PARP1 Monoclonal antibody

Catalog Number: **CL647-66520**

## Basic Information

**Catalog Number:**

CL647-66520

**Size:**

1000 µg/ml

**Source:**

Mouse

**Isotype:**

IgG1

**Immunogen Catalog Number:**

AG19173

**GenBank Accession Number:**

BC037545

**GeneID (NCBI):**

142

**UNIPROT ID:**

P09874

**Full Name:**

poly (ADP-ribose) polymerase 1

**Calculated MW:**

1014 aa, 113 kDa

**Observed MW:**

113-116 kDa, 85-89 kDa

**Purification Method:**

Protein G purification

**CloneNo.:**

1D7D4

**Recommended Dilutions:**

IF 1:50-1:500

**Excitation/Emission maxima  
wavelengths:**

654 nm / 674 nm

## Applications

**Tested Applications:**

FC (Intra), IF/ICC

**Species Specificity:**

Human, mouse, rat

**Positive Controls:**

IF : Neuro-2a cells,

## Background Information

PARP1 (poly(ADP-ribose) polymerase 1) is a nuclear enzyme catalyzing the poly(ADP-ribosyl)ation of many key proteins in vivo. The normal function of PARP1 is the routine repair of DNA damage. Activated by DNA strand breaks, the PARP1 is cleaved into an 85 to 89-kDa COOH-terminal fragment and a 24-kDa NH2-terminal peptide by caspases during the apoptotic process. The appearance of PARP fragments is commonly considered as an important biomarker of apoptosis. In addition to caspases, other proteases like calpains, cathepsins, granzymes and matrix metalloproteinases (MMPs) have also been reported to cleave PARP1 and gave rise to fragments ranging from 42-89-kDa. This antibody was generated against the N-terminal region of human PARP1 and it recognizes the full-length as well as the cleavage of the PARP1.

## Storage

**Storage:**


Store at -20°C. Avoid exposure to light. Stable for one year after shipment.

**Storage Buffer:**


PBS with 50% Glycerol, 0.05% Proclin300, 0.5% BSA, pH 7.3.

Aliquoting is unnecessary for -20°C storage

## Selected Validation Data



Immunofluorescent analysis of (4% PFA) fixed Neuro-2a cells using CoraLite® Plus 647 PARP1 antibody (CL647-66520, Clone: 1D7D4) at dilution of 1:100.



1X10<sup>6</sup> HeLa cells were intracellularly stained with 0.2 ug CoraLite® Plus 647 Anti-Human PARP1 (CL647-66520, Clone:1D7D4) (red), or 0.2 ug Control Antibody. Cells were fixed and permeabilized with Transcription Factor Staining Buffer Kit (PF00011).