

FITC Plus Anti-Human CD107a / LAMP1 (H4A3)

Catalog Number: **FITC-65051**

Basic Information

Catalog Number: FITC-65051	GenBank Accession Number: BC006345	Purification Method: Affinity purification
Size: 100tests, 5 µl/test	GeneID (NCBI): 3916	CloneNo.: H4A3
Source: Mouse	Full Name: lysosomal-associated membrane protein 1	Recommended Dilutions: IF 1:50-1:500
Isotype: IgG1, kappa		Excitation/Emission maxima wavelengths: 494 nm / 520 nm

Applications

Tested Applications: FC, IF	Positive Controls: IF : HeLa cells,
Species Specificity: Human	


Background Information

LAMP1 (CD107a) is a heavily glycosylated membrane protein enriched in the lysosomal membrane. LAMP1 is extensively glycosylated with asparagine-linked oligosaccharides which protect it from intracellular proteolysis (PMID: 10521503). Although LAMP1 is expressed largely in the endosome-lysosomal membrane of cells, it is also found on the plasma membrane (PMID: 16168398). Elevated LAMP1 expression at the cell surface has also been detected during platelet and granulocytic cell activation, as well as in some tumor cells (PMID: 29085473). LAMP1 functions to provide selectins with carbohydrate ligands. This protein has also been shown to be a marker of degranulation on lymphocytes such as CD8+ and NK cells and may also play a role in tumor cell differentiation and metastasis (PMID: 18835598; 29085473; 9426697).


Storage

Storage:
Store at 2-8°C. Avoid exposure to light. Stable for one year after shipment.
Storage Buffer:
PBS with 0.09% sodium azide and 0.5% BSA.


Selected Validation Data



Immunofluorescent analysis of (4% PFA) fixed HeLa cells using FITC-65051 (CD107a antibody) at dilution of 1:50 and CoraLite488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L).



1X10⁶ PHA-activated human PBMCs were surface stained with 5 ul FITC Plus Anti-Human CD107a (FITC-65051, Clone: H4A3). Unactivated PBMCs were used as negative control. Cells were not fixed. Lymphocytes were gated.



1X10⁶ PHA-activated human PBMCs were surface stained with 5 ul FITC Plus Anti-Human CD107a (FITC-65051, Clone: H4A3). Unactivated PBMCs were used as negative control. Cells were not fixed. Lymphocytes were gated.