

## Catalog Number: HZ-1090

## Data Sheet



Animal Component-Free Human cell expressed

Tag-Free E

Endotoxin Free

**ISO** 13485

| Product Description   |  |  |  |  |
|---|--|--|--|--|
| Animal-free Recombinant Human TGF beta 3 is expressed from human 293 cells as a mature disulfide-linked homodimer with apparent molecular mass of 25 kDa. This cytokine is produced in a human cell expression system with serum-free, chemically defined media. The cytokine is greater than 95% pure. |  |  |  |  |
| Alternative Names   | ARVD, FLJ16571, TGF beta 3, TGF beta3, TGFB3                             |  |  |  |
| Source  | Human Embryonic Kidney cells (HEK293). HEK293-derived TGF beta 3 protein |  |  |  |

| Specifications    |   |  |  |  |  |
|-------------------|---|--|--|--|--|
| Test              | Method  | Specification  |  |  |  |
| Activity          | Dose-dependent inhibition of IL-4 induced proliferation of mouse HT-2 cells (BALB/c spleen activated by sheep erythrocytes in the presence of IL-2) | 0.01 - 0.2 ng/mL EC50  |  |  |  |
| Molecular<br>Mass | SDS-PAGE  | 13 kDa reduced, 22 kDa non-reduced,<br>homodimer, non-glycosylated |  |  |  |
| Purity            | SDS-PAGE  | >95%   |  |  |  |
| Endotoxin         | LAL   | <1 EU/µg   |  |  |  |

| Activity Data   | SDS-PAGE   |  |
|---|--|--|
| Recombinant TGF-beta 3   110 100   90 Image: Colspan="2">Image: Colspan="2" Image: Co | Recombinant human TGF<br>beta 3 (HZ-1090) inhibits IL-<br>4 induced proliferation of<br>the HT-2 mouse cell line. HT-<br>2 cells are Balb/c spleen<br>cells activated by sheep<br>erythrocytes in the<br>presence of IL-2. Cell<br>number was quantitatively<br>assessed by PrestoBlue®<br>cell viability reagent. HT-2<br>cells were treated with<br>increasing concentrations of<br>recombinant TGF beta 3 for<br>72 hours. The EC50 was | Reducing Non-Reducing   225 170   130 93   93 70   53 42   30 23   18 14   10 10 |

www.ptglab.com

Document #: FR-QA118-101 Rev 0 Data Sheet Version #: 1 Proteintech Group, Inc. 5500 Pearl Street, Suite 400 Rosemont, IL 60612 t: 1-888-478-4522; f: 1-312-455-8408 Email: proteintech@ptglab.com

| Preparation  |   |  |  |  |
|--|---|--|--|--|
| Shipping<br>Temperature  | ambient temperature   |  |  |  |
| Formulation 50 mM NaOAc pH 3.7 + 1% CHAPS, See Certificate of Analysis for details |   |  |  |  |
| Reconstitution   | Briefly centrifuge the vial before opening. It is recommended to reconstitute the protein to 0.1mg/mL in sterile 4 mM HCl containing 0.1% endotoxin-free recombinant human serum albumin (HSA). |  |  |  |

|                       | Product Form              | Temperature Conditions             | Storage Time<br>(From Date of Receipt) |
|-----------------------|---------------------------|------------------------------------|--|
|                       | Lyophilized               | -20°C to -80°C                     | Until Expiry Date                      |
| Stability and Storage | Lyophilized               | Room Temperature                   | 2 weeks                                |
|                       | Reconstituted as per CofA | -20°C to -80°C                     | 6 months                               |
|                       | Reconstituted as per CofA | 4°C                                | 1 week                                 |
|                       |                           | Avoid repeated freeze-thaw cycles. |  |

www.ptglab.com

Document #: FR-QA118-101 Rev 0 Data Sheet Version #: 1 Proteintech Group, Inc. 5500 Pearl Street, Suite 400 Rosemont, IL 60612 t: 1-888-478-4522; f: 1-312-455-8408 Email: proteintech@ptglab.com