



# IHCeasy Phospho-Histone H3 (Ser10) Ready-To-Use IHC Kit

Catalog Number: KHC1439

**General Information** 

Sample type: FFPE tissue Cited sample type: Reactivity: Human Cited Reactivity:

Assay type: Immunohistochemistry Primary antibody type: Mouse Monoclonal

Secondary antibody type: Polymer-HRP-Goat anti-Mouse

#### Kit Component

| Component                | Size               | Concentration |
|--------------------------|--------------------|---------------|
| Antigen Retrieval Buffer | 100 mL             | 50×           |
| Washing Buffer           | 100 mL ×2          | 20×           |
| Blocking Buffer          | 5 mL               | RTU           |
| Primary Antibody         | 5 mL               | RTU           |
| Secondary Antibody       | 5 mL               | RTU           |
| Chromogen Component A    | 0.2 mL             | RTU           |
| Chromogen Component B    | 4 mL               | RTU           |
| Signal Enhancer          | 5 mL               | RTU           |
| Counter Staining Reagent | 5 mL               | RTU           |
| Mounting Media           | 5 mL               | RTU           |
| Control Slide            | 1 slide (Optional) | FFPE          |
| Datasheet                | 1 Copy             |               |
| Manual                   | 1 Copy             |               |

### **Storage Instructions**

All the reagents are stored at 2-8°C. The kit is stable for 6 months from the date of receipt.


## Background

Phospho-histone-H3 (PHH3) is a core histone protein, which in its phosphorylated state forms the principal constituents of eukaryotic chromatin, with histone H3 being phosphorylated at serine (Ser) 10 or Ser28 as well as its phosphorylation of Ser10 being strongly correlated with the late G2 to M-phase transition in mammalian mitotic cells. On the basis of previous research, a few cell line- and animal model-based researches have displayed an increase in phosphorylation of histone H3 at Ser10 (H3S10ph), the only histone marker that is involved in carcinogenesis and cellular transformation. Histone H3 phosphorylation on serine-10 is specific to mitosis and phosphorylated histone H3 (PHH3) proliferation markers (as counts defined per area or as indices defined per cell numbers) are increasingly being used to evaluate proliferation in various tumors.


#### **Synonyms**

H3 S10, H3 Ser10, H3/A, H3FA, H3S10, HIST1H3A, histone, histone cluster 1, H3a, Histone H3.1, Histone H3/a, PHH3, Phospho-Histone H3 Ser10

## Selected Validation Data



Immunohistochemical analysis of paraffinembedded Jurkat (left) and calyculin A treated Jurkat (right) cells slide using KHC1439 (Phospho-Histone H3 (Ser10) IHC Kit).



Immunohistochemical analysis of paraffinembedded human lung cancer tissue slide using KHC1439 (Phospho-Histone H3 (Ser10) IHC Kit).